arXiv:1106.5090v4 [math.AT] 23 Feb 2012

MORPHISMS OF A_∞-BIALGEBRAS AND APPLICATIONS

SAMSON SANEBLIDZE[1] AND RONALD UMBLE[2]

Abstract. We define the notion of a *relative matrad* and construct a new family of polytopes $JJ = \{JJ_{n,m} = JJ_{m,n}\}_{m,n\geq 1}$, called *bimultiplihedra*, of which $JJ_{n,1} = JJ_{1,n}$ is the multiplihedron J_n. We realize the free relative matrad $r\mathcal{H}_\infty$ by extending the free $\mathcal{A}_\infty$-bimodule structure on cellular chains of multiplihedra to a free $\mathcal{H}_\infty$-bimodule structure on cellular chains $C_*(JJ)$. A *morphism* $G : A \Rightarrow B$ of A_∞-bialgebras is the image of a map $C_*(JJ) \to Hom(TA, TB)$ of relative matrads. We prove that the homology of every A_∞-bialgebra over a commutative ring with unity (in particular, the bialgebra $S_*(\Omega X; \mathbb{Z})$ of singular chains on Moore base pointed loops) admits an induced A_∞-bialgebra structure. We extend the classical Bott-Samelson isomorphism to an isomorphism of A_∞-bialgebras and identify the A_∞-bialgebra structure of $H_*(\Omega\Sigma X; \mathbb{Q})$. This is the first known nontrivial rational homology invariant for $\Omega\Sigma X$. For each $n \geq 2$, we construct a space X_n whose loop cohomology $H = H^*(\Omega X_n; \mathbb{Z}_2)$ comes equipped with a nontrivial operation $\omega_2^n : H \otimes H \to H^{\otimes n}$. To our knowledge, ω_2^n is the first known example of a non-operadic operation on loop cohomology.

1. Introduction

In [21] and [22] we constructed the objects in the category of A_∞-bialgebras; in this paper we construct the morphisms. Given a module H over a (graded or ungraded) commutative ring R with unity, consider the bigraded module $M = \{M_{n,m} = Hom(H^{\otimes m}, H^{\otimes n})\}$, the bisequence submodule $\mathbf{M} \subseteq TTM$ (the tensor module of TM), and the canonical upsilon product $\Upsilon : \mathbf{M} \times \mathbf{M} \to \mathbf{M}$. Given an arbitrary family of R-multilinear operations

$$\omega = \{\omega_m^n \in M_{n,m} \mid |\omega_m^n| = m + n - 3\}_{m,n\geq 1},$$

the sum $\sum \omega_m^n$ extends uniquely to its biderivative $d_\omega \in \mathbf{M}$, and there is a (non-bilinear) product $\circledcirc : M \times M \overset{d_-\times d_-}{\longrightarrow} \mathbf{M} \times \mathbf{M} \overset{\Upsilon}{\rightarrow} \mathbf{M} \overset{\text{proj}}{\rightarrow} M$, which extends Gerstenhaber's $\circ_i$-products on $Hom(TH, H) \oplus Hom(H, TH)$ [6]. The family ω defines an A_∞-bialgebra structure on H whenever $\omega \circledcirc \omega = 0$, and the structure relations appear as bihomogeneous components of this equation. Indeed, the structure relation in bidegree (m, n) is modeled on the biassociahedron $KK_{n,m}$, and the cellular chains $C_*(KK)$ realize the free matrad $\mathcal{H}_\infty$. Thus (H, ω) is an A_∞-bialgebra if ω is the image of a map $C_*(KK) \to M$ of matrads, i.e., H is an algebra over $\mathcal{H}_\infty$.

Date: February 22, 2012.

1991 *Mathematics Subject Classification.* Primary 55P35, 55P99 ; Secondary 52B05.

Key words and phrases. A_∞-bialgebra, biassociahedron, bimultiplihedron, loop space, matrad, operad, permutahedron, relative matrad .

[1] This research described in this publication was made possible in part by Award No. GM1-2083 of the U.S. Civilian Research and Development Foundation for the Independent States of the Former Soviet Union (CRDF) and by Award No. 99-00817 of INTAS.

[2] This research funded in part by a Millersville University faculty research grant.

In this paper we define the notion of a *relative matrad* and construct a new family of polytopes $JJ = \{JJ_{n,m} = JJ_{m,n}\}_{m,n\geq 1}$, called *bimultiplihedra*, of which $JJ_{n,1} = JJ_{1,n}$ is the multiplihedron J_n. We realize the free relative matrad $r\mathcal{H}_\infty$ by extending the free $\mathcal{A}_\infty$-bimodule structure on cellular chains of multiplihedra to a free $\mathcal{H}_\infty$-bimodule structure on cellular chains $C_*(JJ)$. On the other hand, given A_∞-bialgebras A and B, we consider $Hom(TA, TB)$ as an $\mathcal{E}nd_{TB}$-$\mathcal{E}nd_{TA}$-bimodule and define a *morphism* $G : A \Rightarrow B$ as the image of a map $C_*(JJ) \to Hom(TA, TB)$ of relative matrads, i.e., G is a bimodule over $\mathcal{H}_\infty$.

Given an A_∞-bialgebra B and a homology isomorphism $g : H_*(B) \to B$, we apply our generalization of the Basic Perturbation Lemma to prove that under mild conditions, the A_∞-bialgebra structure on B pulls back along g to an A_∞-bialgebra structure on $H_*(B)$, and any two such structures so obtained are isomorphic. This is a special case of our main result:

Theorem 2. *Let B be an A_∞-bialgebra with homology $H = H_*(B)$, let (RH, d) be a free R-module resolution of H, and let h be a perturbation of d such that $g : (RH, d + h) \to (B, d_B)$ is a homology isomorphism. Then*

- *(i)* (Existence) *g induces an A_∞-bialgebra structure ω_{RH} on RH and extends to a map $G : RH \Rightarrow B$ of A_∞-bialgebras.*
- *(ii)* (Uniqueness) *(ω_{RH}, G) is unique up to isomorphism.*

In particular, given a topological space X and a field F, the bialgebra structure on simplicial singular chains $S_*(\Omega X; F)$ of Moore base pointed loops pulls back to an A_∞-bialgebra structure on $H_*(\Omega X; F)$ and the induced A_∞-(co)algebra substructures are exactly the ones observed earlier by Gugenheim [8] and Kadeishvili [11]. Furthermore, we show that the A_∞-coalgebra structure on $H_*(X; F)$ extends to an A_∞-bialgebra structure on the tensor algebra $T^a\tilde{H}_*(X; F)$, which is trivial if and only if the A_∞-coalgebra structure on $H_*(X; F)$ is trivial. We extend the classical Bott-Samelson Isomorphism $t_* : T^a\tilde{H}_*(X; F) \overset{\approx}{\to} H_*(\Omega\Sigma X; F)$ to an isomorphism of A_∞-bialgebras and conclude that the A_∞-bialgebra structure of $H_*(\Omega\Sigma X; F)$ is trivial if and only if the A_∞-coalgebra structure on $H_*(X; F)$ is trivial (Theorem 4). This is the A_∞-bialgebra structure of $H_*(\Omega\Sigma X; \mathbb{Q})$ is the first nontrivial rational homology invariant for $\Omega\Sigma X$ (Corollary 2). Finally, for each $n \geq 2$, we construct a space X_n and a nontrivial operation $\omega_2^n : H^*(\Omega X_n; \mathbb{Z}_2)^{\otimes 2} \to H^*(\Omega X_n; \mathbb{Z}_2)^{\otimes n}$ defined in terms of the action of the Steenrod algebra $\mathcal{A}_2$ on $H^*(X_n; \mathbb{Z}_2)$. To our knowledge, ω_2^n is the first known example of a non-operadic operation on cohomology.

The paper is organized as follows: Section 2 reviews the theory of matrads and the related notation used throughout the paper; see [22] for details. We construct relative matrads in Section 3 and the bimultiplihedra in Section 4. We define morphisms of A_∞-bialgebras in Section 5. We prove our generalization of the classical Basic Perturbation Lemma (Theorem 1) and our main result (Theorem 2) in Section 6. We conclude with various applications and examples in Section 7.

2. MATRADS

Let $M = \{M_{n,m}\}_{m,n\geq 1}$ be a bigraded module over a (graded or ungraded) commutative ring R with unity 1_R. In particular, given a graded R-module H, we set $M_{n,m} = Hom(H^{\otimes m}, H^{\otimes n})$ and think of $\alpha_m^n \in M_{n,m}$ as some composition of multilinear operations θ_i^j pictured as a possibly non-planar upward-directed *graph* with m inputs and n outputs.

Each pair of matrices $X = [x_{ij}]\,, Y = [y_{ij}] \in \mathbb{N}^{q\times p}$, $p, q \geq 1$, uniquely determines a submodule

$$M_{Y,X} = \left(M_{y_{11},x_{11}} \otimes \cdots \otimes M_{y_{1p},x_{1p}}\right) \otimes \cdots \otimes \left(M_{y_{q1},x_{q1}} \otimes \cdots \otimes M_{y_{qp},x_{qp}}\right) \subset TTM.$$

Fix a set of bihomogeneous module generators $G \subset M$. A *monomial in* TM is an element of $G^{\otimes p}$ and a *monomial in* TTM is an element of $\left(G^{\otimes p}\right)^{\otimes q}$. Thus $A \in \left(G^{\otimes p}\right)^{\otimes q}$ is naturally represented by the $q \times p$ matrix

$$[A] = \left[\begin{array}{ccc} \alpha_{x_{1,1}}^{y_{1,1}} & \cdots & \alpha_{x_{1,p}}^{y_{1,p}} \\ \vdots & & \vdots \\ \alpha_{x_{q,1}}^{y_{q,1}} & \cdots & \alpha_{x_{q,p}}^{y_{q,p}} \end{array}\right]$$

with entries in G and rows identified with elements of $G^{\otimes p}$. We refer to A as a $q \times p$ *monomial* (we use the symbols A and $[A]$ interchangeably). Consequently,

$$\left(M^{\otimes p}\right)^{\otimes q} = \bigoplus_{X,Y \in \mathbb{N}^{q\times p}} M_{Y,X}$$

and we refer to

$$\overline{\mathbf{M}} = \bigoplus_{\substack{X,Y \in \mathbb{N}^{q\times p} \\ p,q \geq 1}} M_{Y,X} \quad \text{and} \quad \overline{\mathbf{V}} = \bigoplus_{\substack{X,Y \in \mathbb{N}^{1\times p} \cup \mathbb{N}^{q\times 1} \\ p,q \geq 1}} M_{Y,X}$$

as the *matrix* and *vector* submodules of TTM, respectively. The matrix transpose $A \mapsto A^T$ induces the permutation of tensor factors $\sigma_{p,q} : \left(M^{\otimes p}\right)^{\otimes q} \overset{\approx}{\to} \left(M^{\otimes q}\right)^{\otimes p}$ given by

$$\begin{gathered} \left(\alpha_{x_{1,1}}^{y_{1,1}} \otimes \cdots \otimes \alpha_{x_{1,p}}^{y_{1,p}}\right) \otimes \cdots \otimes \left(\alpha_{x_{q,1}}^{y_{q,1}} \otimes \cdots \otimes \alpha_{x_{q,p}}^{y_{q,p}}\right) \mapsto \\ \left(\alpha_{x_{1,1}}^{y_{1,1}} \otimes \cdots \otimes \alpha_{x_{q,1}}^{y_{q,1}}\right) \otimes \cdots \otimes \left(\alpha_{x_{1,p}}^{y_{1,p}} \otimes \cdots \otimes \alpha_{x_{q,p}}^{y_{q,p}}\right). \end{gathered}$$

Let $\bar{X}$ and $\bar{Y}$ denote matrices in $\mathbb{N}^{q\times p}$ with constant columns $(x_i)^T$ and constant rows (y_j), respectively, and consider $q \times p$ monomials

$$A = \left[\begin{array}{ccc} \alpha_{x_1}^{y_{1,1}} & \cdots & \alpha_{x_p}^{y_{1,p}} \\ \vdots & & \vdots \\ \alpha_{x_1}^{y_{q,1}} & \cdots & \alpha_{x_p}^{y_{q,p}} \end{array}\right] \in M_{Y,\bar{X}} \quad \text{and} \quad B = \left[\begin{array}{ccc} \beta_{x_{1,1}}^{y_1} & \cdots & \beta_{x_{1,p}}^{y_1} \\ \vdots & & \vdots \\ \beta_{x_{q,1}}^{y_q} & \cdots & \beta_{x_{q,p}}^{y_q} \end{array}\right] \in M_{\bar{Y},X}.$$

The *row leaf sequence of* A is the p-tuple $\operatorname{rls}(A) = (x_1, \ldots, x_p)$; the *column leaf sequence of* B is the q-tuple $\operatorname{cls}(B) = (y_1, \ldots, y_q)^T$. Define

$$\overline{\mathbf{M}}_{\text{row}} = \bigoplus_{\bar{\mathbf{X}},Y \in \mathbb{N}^{q\times p};\ p,q \geq 1} M_{Y,\bar{X}} \quad \text{and} \quad \overline{\mathbf{M}}^{\text{col}} = \bigoplus_{X,\bar{Y} \in \mathbb{N}^{q\times p};\ p,q \geq 1} M_{\bar{Y},X};$$

then

$$\mathbf{M} = \overline{\mathbf{M}}_{\text{row}} \cap \overline{\mathbf{M}}^{\text{col}}$$

is the *bisequence submodule* of TTM, and a $q \times p$ monomial $A \in \mathbf{M}$ is represented as a *bisequence matrix*

$$A = \left[\begin{array}{ccc} \alpha_{x_1}^{y_1} & \cdots & \alpha_{x_p}^{y_1} \\ \vdots & & \vdots \\ \alpha_{x_1}^{y_q} & \cdots & \alpha_{x_p}^{y_q} \end{array}\right].$$

Unless otherwise stated, we shall assume that $\mathbf{x}\times\mathbf{y}\in(x_1,\ldots,x_p)\times(y_1,\ldots,y_q)^T\in\mathbb{N}^{1\times p}\times\mathbb{N}^{q\times 1}$, $p,q\geq 1$; we will often express $\mathbf{y}$ as a row vector. Let

$$\mathbf{M}_{\mathbf{x}}^{\mathbf{y}}=\left\langle A\in\mathbf{M}\mid\mathbf{x}=\operatorname{rls}(A)\ \text{ and }\mathbf{y}=\operatorname{cls}(A)\right\rangle;$$

then

$$\mathbf{M}=\bigoplus_{\mathbf{x}\times\mathbf{y}}\mathbf{M}_{\mathbf{x}}^{\mathbf{y}}.$$

By identifying $\left(H^{\otimes q}\right)^{\otimes p}\approx\left(H^{\otimes p}\right)^{\otimes q}$ with $(q,p)\in\mathbb{N}^2$, we think of a $q\times p$ monomial $A\in\mathbf{M}_{\mathbf{x}}^{\mathbf{y}}$ as an operator on the positive integer lattice $\mathbb{N}^2$, pictured as an arrow $(|\mathbf{x}|,q)\mapsto(p,|\mathbf{y}|)$, where $|\mathbf{u}|=\Sigma u_i$; but unfortunately, this representation is not faithful. The *bisequence vector submodule* is the intersection

$$\mathbf{V}=\overline{\mathbf{V}}\cap\mathbf{M}=\bigoplus_{s,t\in\mathbb{N}}\mathbf{M}_s^{\mathbf{y}}\oplus\mathbf{M}_{\mathbf{x}}^t.$$

A submodule

$$\mathbf{W}=M\oplus\bigoplus_{\mathbf{x},\mathbf{y}\notin\mathbb{N};\ s,t\in\mathbb{N}}\mathbf{W}_s^{\mathbf{y}}\oplus\mathbf{W}_{\mathbf{x}}^t\subseteq\mathbf{V}$$

is *telescoping* if for all $\mathbf{x},\mathbf{y},s,t$

(i) $\mathbf{W}_s^{\mathbf{y}}\subseteq\mathbf{M}_s^{\mathbf{y}}$ and $\mathbf{W}_{\mathbf{x}}^t\subseteq\mathbf{M}_{\mathbf{x}}^t$;

(ii) $\alpha_s^{y_1}\otimes\cdots\otimes\alpha_s^{y_q}\in\mathbf{W}_s^{\mathbf{y}}$ implies $\alpha_s^{y_1}\otimes\cdots\otimes\alpha_s^{y_j}\in\mathbf{W}_s^{y_1\cdots y_j}$ for all $j<q$;

(iii) $\beta_{x_1}^t\otimes\cdots\otimes\beta_{x_p}^t\in\mathbf{W}_{\mathbf{x}}^t$ implies $\beta_{x_1}^t\otimes\cdots\otimes\beta_{x_i}^t\in\mathbf{W}_{x_1\cdots x_i}^t$ for all $i<p$.

Thus the truncation maps $\tau:\mathbf{W}_s^{y_1\cdots y_j}\to\mathbf{W}_s^{y_1\cdots y_{j-1}}$ and $\tau:\mathbf{W}_{x_1\cdots x_i}^t\to\mathbf{W}_{x_1\cdots x_{i-1}}^t$ determine the following "telescoping" sequences of submodules:

$$\tau\left(\mathbf{W}_s^{\mathbf{y}}\right)\subseteq\tau^2\left(\mathbf{W}_s^{\mathbf{y}}\right)\subseteq\cdots\subseteq\tau^{q-1}\left(\mathbf{W}_s^{\mathbf{y}}\right)=\mathbf{W}_s^{y_1}$$
$$\tau\left(\mathbf{W}_{\mathbf{x}}^t\right)\subseteq\tau^2\left(\mathbf{W}_{\mathbf{x}}^t\right)\subseteq\cdots\subseteq\tau^{p-1}\left(\mathbf{W}_{\mathbf{x}}^t\right)=\mathbf{W}_{x_1}^t.$$

In general, $\mathbf{W}_{\mathbf{x}}^t$ is an *additive* submodule of $\mathbf{M}_{x_1}^t\otimes\cdots\otimes\mathbf{M}_{x_p}^t$ and does *not* necessarily decompose as $B_1\otimes\cdots\otimes B_p$ with $B_i\subseteq\mathbf{M}_{x_i}^t$.

The *telescopic extension* of a telescoping submodule $\mathbf{W}\subseteq\mathbf{V}$ is the submodule of matrices $\mathcal{W}\subseteq\overline{\mathbf{M}}$ with the following properties: If $A=\left[\alpha_{x_{i,j}}^{y_{i,j}}\right]$ is a $q\times p$ monomial in $\mathcal{W}$ and

(i) $\left[\alpha_{x_{i,j}}^{y_{i,j}}\ \cdots\ \alpha_{x_{i,j+m}}^{y_{i,j+m}}\right]$ is a string in the i^{th} row of A such that $y_{i,j}=\cdots=y_{i,j+m}=t$, then $\alpha_{x_{i,j}}^t\otimes\cdots\otimes\alpha_{x_{i,j+m}}^t\in\mathbf{W}_{x_{i,j},\ldots,x_{i,j+m}}^t$.

(ii) $\left[\alpha_{x_{i,j}}^{y_{i,j}}\ \cdots\ \alpha_{x_{i+r,j}}^{y_{i+r,j}}\right]^T$ is a string in the j^{th} column of A such that $x_{i,j}=\cdots=x_{i+r,j}=s$, then $\alpha_s^{y_{i,j}}\otimes\cdots\otimes\alpha_s^{y_{i+r,j}}\in\mathbf{W}_s^{y_{i,j},\ldots,y_{i+r,j}}$.

Thus if $A\in\mathbf{M}_{\mathbf{x}}^{\mathbf{y}}\cap\mathcal{W}$, the i^{th} row of A lies in $\mathbf{W}_{\mathbf{x}}^{y_i}$ and the j^{th} column of A lies in $\mathbf{W}_{x_j}^{\mathbf{y}}$.

Definition 1. *A pair* $A\otimes B=\left[\alpha_{v_{kl}}^{y_{kl}}\right]\otimes\left[\beta_{x_{ij}}^{u_{ij}}\right]$ *of* $(q\times s,t\times p)$ *monomials in* $\overline{\mathbf{M}}\otimes\overline{\mathbf{M}}$ *is a*

(i) ***Transverse Pair*** *(TP) if* $s=t=1$, $u_{1,j}=q$, *and* $v_{k,1}=p$ *for all* j *and* k, *i.e., setting* $x_j=x_{1,j}$ *and* $y_k=y_{k,1}$ *gives*

$$A\otimes B=\begin{bmatrix}\alpha_p^{y_1}\\ \vdots\\ \alpha_p^{y_q}\end{bmatrix}\otimes\begin{bmatrix}\beta_{x_1}^q & \cdots & \beta_{x_p}^q\end{bmatrix}\in\mathbf{M}_p^{\mathbf{y}}\otimes\mathbf{M}_{\mathbf{x}}^q.$$

(ii) ***Block Transverse Pair*** *(BTP) if there exist $t \times s$ block decompositions $A = \left[A'_{k',l}\right]$ and $B = \left[B'_{i,j'}\right]$ such that $A'_{il} \otimes B'_{il}$ is a TP for all i and l.*

A pair $A \otimes B \in \mathbf{M}^{\mathbf{y}}_{\mathbf{v}} \otimes \mathbf{M}^{\mathbf{u}}_{\mathbf{x}}$ is a BTP if and only if $\mathbf{x} \times \mathbf{y} \in \mathbb{N}^{1\times|\mathbf{v}|} \times \mathbb{N}^{|\mathbf{u}|\times 1}$ if and only if the initial point of arrow A coincides with the terminal point of arrow B as operators on $\mathbb{N}^2$.

Given a family of maps $\bar{\gamma} = \{M^{\otimes q} \otimes M^{\otimes p} \to M\}_{p,q\geq 1}$, let $\gamma = \{\gamma^{\mathbf{y}}_{\mathbf{x}} : \mathbf{M}^{\mathbf{y}}_{p} \otimes \mathbf{M}^{q}_{\mathbf{x}} \to \mathbf{M}^{|\mathbf{y}|}_{|\mathbf{x}|}\}$. Then γ induces a global product $\Upsilon : \overline{\mathbf{M}} \otimes \overline{\mathbf{M}} \to \overline{\mathbf{M}}$ defined by

$$\Upsilon\left(A \otimes B\right) = \begin{cases} \left[\gamma\left(A'_{ij} \otimes B'_{ij}\right)\right], & A \otimes B \text{ is a BTP} \\ 0, & \text{otherwise,} \end{cases} \tag{2.1}$$

where $A'_{ij} \otimes B'_{ij}$ is the $(i,j)^{th}$ TP in the BTP decomposition of $A \otimes B$. Obviously Υ is closed in both $\overline{\mathbf{M}}_{\text{row}}$ and $\overline{\mathbf{M}}^{\text{col}}$; consequently, Υ is closed in $\mathbf{M}$. We denote the Υ-product by "$\cdot$" or juxtaposition; when $A \otimes B = \left[\alpha^{y_j}_{p}\right]^T \otimes \left[\beta^{q}_{x_i}\right]$ is a TP, we write

$$AB = \gamma(\alpha^{y_1}_{p}, \ldots, \alpha^{y_q}_{p}; \beta^{q}_{x_1}, \ldots, \beta^{q}_{x_p}).$$

As an arrow, AB "transgresses" from the x-axis to the y-axis $\mathbb{N}^2$.

Let $\mathbf{1}^{1\times p} = (1, \ldots, 1) \in \mathbb{N}^{1\times p}$ and $\mathbf{1}^{q\times 1} = (1, \ldots, 1)^T \in \mathbb{N}^{q\times 1}$; we often suppress the exponents when the context is clear.

Definition 2. *A* ***prematrad*** *(M, γ, η) is a bigraded R-module $M = \{M_{n,m}\}_{m,n\geq 1}$ together with a family of structure maps $\gamma = \{\gamma^{\mathbf{y}}_{\mathbf{x}} : \mathbf{M}^{\mathbf{y}}_{p} \otimes \mathbf{M}^{q}_{\mathbf{x}} \to \mathbf{M}^{|\mathbf{y}|}_{|\mathbf{x}|}\}$ and a unit $\eta : R \to \mathbf{M}^1_1$ such that*

- *(i)* *$\Upsilon\left(\Upsilon\left(A; B\right); C\right) = \Upsilon\left(A; \Upsilon\left(B; C\right)\right)$ whenever $A \otimes B$ and $B \otimes C$ are BTPs in $\overline{\mathbf{M}} \otimes \overline{\mathbf{M}}$;*
- *(ii)* *the following compositions are the canonical isomorphisms:*

$$R^{\otimes b} \otimes \mathbf{M}^b_a \overset{\eta^{\otimes b}\otimes \mathrm{Id}}{\longrightarrow} \mathbf{M}^{\mathbf{1}^{b\times 1}}_1 \otimes \mathbf{M}^b_a \overset{\gamma^{\mathbf{1}^{b\times 1}}_a}{\longrightarrow} \mathbf{M}^b_a;$$

$$\mathbf{M}^b_a \otimes R^{\otimes a} \overset{\mathrm{Id}\otimes \eta^{\otimes a}}{\longrightarrow} \mathbf{M}^b_a \otimes \mathbf{M}^1_{\mathbf{1}^{1\times a}} \overset{\gamma^b_{\mathbf{1}^{1\times a}}}{\longrightarrow} \mathbf{M}^b_a.$$

We denote the element $\eta(1_R)$ by $\mathbf{1_M}$.

A ***morphism of prematrads*** *(M, γ) and (M', γ') is a map $f : M \to M'$ such that $f\gamma^{\mathbf{y}}_{\mathbf{x}} = {\gamma'}^{\mathbf{y}}_{\mathbf{x}}(f^{\otimes q} \otimes f^{\otimes p})$ for all $\mathbf{x} \times \mathbf{y}$.*

Although Υ fails to act associatively on $\overline{\mathbf{M}}$, Axiom (i) implies that $(AB)\,C = A\,(BC)$ whenever $A, B, C \in \overline{\mathbf{M}}$, $AB \neq 0$, and $BC \neq 0$. On the other hand, Υ acts associatively on $\mathbf{M}$. Given a bisequence matrix $A^{q\times p} \in \mathbf{M}^{\mathbf{y}}_{\mathbf{x}}$, consider the constant (bisequence) matrices $\mathbf{1}^{q\times|\mathbf{x}|}$ and $\mathbf{1}^{|\mathbf{y}|\times p}$ whose entries are constantly $\mathbf{1_M}$. Then $\Upsilon\left(\mathbf{1}^{|\mathbf{y}|\times p}; A\right) = A = \Upsilon\left(A; \mathbf{1}^{q\times|\mathbf{x}|}\right)$ by Axiom (ii).

Definition 3. *Let (M, γ, η) be a prematrad. A string of matrices $A_s \cdots A_1$ is* ***basic in*** *$\mathbf{M}^n_m$ if*

- *(i)* *$A_1 \in \mathbf{M}^b_{\mathbf{x}}$, $|\mathbf{x}| = m$,*
- *(ii)* *$A_i \in \overline{\mathbf{M}} \smallsetminus \{\mathbf{1}^{q\times p} \mid p, q \in \mathbb{N}\}$ for all i,*
- *(iii)* *$A_s \in \mathbf{M}^{\mathbf{y}}_a$, $|\mathbf{y}| = n$, and*
- *(iv)* *some association of $A_s \cdots A_1$ defines a sequence of BTPs and non-zero Υ-products.*

Our next definition constructs a bigraded set $G^{\mathrm{pre}} = G^{\mathrm{pre}}_{*,*}$ in which $G^{\mathrm{pre}}_{n,m}$ is defined in terms of the intermediate set

$$G_{[n,m]} = \bigcup_{\substack{i\leq m,\ j\leq n,\\ i+j<m+n}} G^{\mathrm{pre}}_{j,i}\,.$$

We denote the set of matrices over $G_{[n,m]}$ by $\overline{\mathbf{G}}_{[n,m]}$, the set of matrices over G^{pre} by $\overline{\mathbf{G}}$, and the subset of bisequence matrices in $\overline{\mathbf{G}}$ by $\mathbf{G}$.

Definition 4. *Let $\Theta = \left\langle \theta^n_m \mid \theta^1_1 = \mathbf{1}\right\rangle_{m,n\geq 1}$ be a free bigraded R-module generated by singletons θ^n_m and set $G^{\mathrm{pre}}_{1,1} = \mathbf{1}$. Inductively, if $m+n \geq 3$ and $G^{\mathrm{pre}}_{j,i}$ has been constructed for $i \leq m$, $j \leq n$, and $i+j < m+n$, define*

$$G^{\mathrm{pre}}_{n,m} = \theta^n_m \cup \left\{\textit{basic strings } A_s \cdots A_1 \textit{ in } \mathbf{G}^n_m \textit{ with } s \geq 2\right\}.$$

*Let $\sim$ be the equivalence relation on $G^{\mathrm{pre}} = G^{\mathrm{pre}}_{*,*}$ generated by $[A_{ij}B_{ij}] \sim [A_{ij}]\,[B_{ij}]$ if and only if $[A_{ij}] \times [B_{ij}] \in \overline{\mathbf{G}} \times \overline{\mathbf{G}}$ is a BTP, and let $F^{\mathrm{pre}}(\Theta) = \langle G^{\mathrm{pre}} \diagup \sim\rangle$. The **free prematrad generated by** Θ is the prematrad $(F^{\mathrm{pre}}(\Theta), \gamma, \eta)$, where γ is juxtaposition and $\eta : R \to F^{\mathrm{pre}}_{1,1}(\Theta)$ is given by $\eta(1_R) = \mathbf{1}$.*

Let $\mathcal{W}$ be the telescopic extension of a telescoping submodule $\mathbf{W}$. If $A \otimes B$ is a BTP in $\mathcal{W} \otimes \mathcal{W}$, each TP $A' \otimes B'$ in $A \otimes B$ lies in $\mathbf{W}^{\mathbf{y}}_p \otimes \mathbf{W}^q_{\mathbf{x}}$ for some $\mathbf{x}, \mathbf{y}, p, q$. Consequently, $\gamma_{\mathbf{W}}$ extends to a global product $\Upsilon : \mathcal{W} \otimes \mathcal{W} \to \mathcal{W}$ as in (2.1). In fact, $\mathcal{W}$ is the smallest matrix submodule containing $\mathbf{W}$ on which Υ is well-defined.

Definition 5. *Let $\mathbf{W}$ be a telescoping submodule of TTM and let $\mathcal{W}$ be its telescopic extension. Let $\gamma_{\mathbf{W}} = \left\{\gamma^{\mathbf{y}}_{\mathbf{x}} : \mathbf{W}^{\mathbf{y}}_p \otimes \mathbf{W}^q_{\mathbf{x}} \to \mathbf{W}^{|\mathbf{y}|}_{|\mathbf{x}|}\right\}$ be a structure map and let $\eta : R \to \mathbf{M}$. The triple $(M, \gamma_{\mathbf{W}}, \eta)$ is a **local prematrad (with domain $\mathbf{W}$)** if the following axioms are satisfied:*

- *(i)* $\mathbf{W}^1_{\mathbf{x}} = \mathbf{M}^1_{\mathbf{x}}$ *and* $\mathbf{W}^{\mathbf{y}}_1 = \mathbf{M}^{\mathbf{y}}_1$ *for all* $\mathbf{x}, \mathbf{y}$*;*
- *(ii)* $\Upsilon(\Upsilon(A;B);C) = \Upsilon(A;\Upsilon(B;C))$ *whenever $A \otimes B$ and $B \otimes C$ are BTPs in $\mathcal{W} \otimes \mathcal{W}$;*
- *(iii) the prematrad unit axiom holds for $\gamma_{\mathbf{W}}$.*

Recall that each codimension k face of the permutahedron P_m is identified with two planar rooted trees with levels (PLTs)–an up-rooted PLT with $m+1$ leaves and $k+1$ levels, and its down-rooted mirror image (see [14], [20]). Define the $(m,1)$-*row descent sequence* of the m-leaf up-rooted corolla $\curlywedge_m$ to be $\mathbf{m} = (m)$. Given an up-rooted PLT T with $k \geq 2$ levels, successively remove the levels of T and obtain a sequence of subtrees $T = T^k, T^{k-1}, \ldots, T^1$, in which $T^i - T^{i-1}$ is the sequence of corollas $\curlywedge_{m_{i,1}} \cdots \curlywedge_{m_{i,r_i}}$. Recover T inductively by attaching $\curlywedge_{m_{i,j}}$ to the j^{th} leaf of T^{i-1}. Define the i^{th} *leaf sequence* of T to be the row matrix $\mathbf{m}_i = (m_{i,1}, \ldots, m_{i,r_i})$ and the (m,k)-*row descent sequence of* T to be the k-tuple of row matrices $(\mathbf{m}_1, \ldots, \mathbf{m}_k)$. Dually, define the $(n,1)$-*column descent sequence* of the n-leaf down-rooted corolla $\curlyvee^n$ to be $\mathbf{n} = (n)$. Define the $(n,1)$-*column descent sequence* of the n-leaf down-rooted corolla $\curlywedge_m$ to be $\mathbf{m} = (m)$. Given a down-rooted PLT T with $l \geq 2$ levels, successively remove the levels of T and obtain a sequence of subtrees $T = T^l, T^{l-1}, \ldots, T^1$, in which $T^i - T^{i-1}$ is the sequence of corollas $\curlyvee^{n_{i,1}} \cdots \curlyvee^{n_{i,s_i}}$. Recover T inductively by attaching $\curlyvee^{n_{i,j}}$ to the j^{th} leaf of T^{i-1} for each i. Define the i^{th} *leaf sequence* of T to be the column matrix $\mathbf{n}_i = (n_{i,1}, \ldots, n_{i,s_i})^T$ and the (n,l)-*column descent sequence of* T to be the l-tuple of column matrices $(\mathbf{n}_l, \ldots, \mathbf{n}_1)$.

Let $\mathcal{W}_{\text{row}} = \mathcal{W} \cap \overline{\mathbf{M}}_{\text{row}}$ and $\mathcal{W}^{\text{col}} = \mathcal{W} \cap \overline{\mathbf{M}}^{\text{col}}$.

Definition 6. *Given a local prematrad* $(M, \gamma_{\mathbf{W}})$ *with domain* $\mathbf{W}$, *let* $\zeta \in M_{*,m}$ *and* $\xi \in M_{n,*}$ *be elements with* $m, n \geq 2$.

(i) *A* ***row factorization of*** ζ ***with respect to*** $\mathbf{W}$ *is a* Υ*-factorization* $A_1 \cdots A_k = \zeta$ *such that* $A_j \in \mathcal{W}_{\text{row}}$ *and* $\operatorname{rls}(A_j) \neq \mathbf{1}$ *for all* j. *The sequence* $\alpha = (\operatorname{rls}(A_1), \ \ldots, \operatorname{rls}(A_k))$ *is the related* (m,k)*-row* ***descent sequence of*** ζ.

(ii) *A* ***column factorization of*** ξ ***with respect to*** $\mathbf{W}$ *is a* Υ*-factorization* $B_l \cdots B_1 = \xi$ *such that* $B_i \in \mathcal{W}^{\text{col}}$ *and* $\operatorname{cls}(B_i) \neq \mathbf{1}$ *for all* i. *The sequence* $\beta = (\operatorname{cls}(B_l), \ \ldots, \operatorname{cls}(B_1))$ *is the related* (n,l)*-**column descent sequence of*** ξ.

Given a local prematrad $(M, \gamma_{\mathbf{W}})$ and elements $A \in M_{*,s}$ and $B \in M_{t,*}$ with $s, t \geq 2$, choose a row factorization $A_1 \cdots A_k$ of A with respect to $\mathbf{W}$ having (s,k)-row descent sequence α, and a column factorization $B_l \cdots B_1$ of B with respect to $\mathbf{W}$ having (t,l)-column descent sequence β. Then α identifies A with an up-rooted s-leaf, k-level PLT and a codimension $k-1$ face $\hat{e}_A$ of P_{s-1}, and β identifies B with a down-rooted t-leaf, l-level PLT and a codimension $l-1$ face $\check{e}_B$ of P_{t-1}. Extending to Cartesian products, identify the monomials $A = A_1 \otimes \cdots \otimes A_q \in (M_{*,s})^{\otimes q}$ and $B = B_1 \otimes \cdots \otimes B_p \in (M_{t,*})^{\otimes p}$ with the product cells

$$\hat{e}_A = \hat{e}_{A_1} \times \cdots \times \hat{e}_{A_q} \subset P_{s-1}^{\times q} \quad \text{and} \quad \overset{\vee}{e}_B = \check{e}_{B_1} \times \cdots \times \check{e}_{B_p} \subset P_{t-1}^{\times p}.$$

Now consider the S-U diagonal Δ_P and recall that there is a k-subdivision $P_r^{(k)}$ of P_r and a cellular inclusion $P_r^{(k)} \hookrightarrow \Delta^{(k)}(P_r) \subset P_r^{\times k+1}$ for each k and r (see [22]). Thus for each $q \geq 2$, the product cell $\hat{e}_A$ either *is* or *is not* a subcomplex of $\Delta^{(q-1)}(P_{s-1}) \subset P_{s-1}^{\times q}$, and dually for $\overset{\vee}{e}_B$. Let $\mathbf{x}_{m,i}^{p} = (1, \ldots, m, \ldots, 1) \in \mathbb{N}^{1 \times p}$ with m in the i^{th} position and let $\mathbf{y}_q^{n,j} = (1, \ldots, n, \ldots, 1)^T \in \mathbb{N}^{q \times 1}$ with n in the j^{th} position.

Definition 7. *The* ***(left) configuration module*** *of a local prematrad* $(M, \gamma_{\mathbf{W}})$ *is the* R*-module*

$$\Gamma(M, \gamma_{\mathbf{W}}) = M \oplus \bigoplus_{\mathbf{x}, \mathbf{y} \notin \mathbb{N};\ s,t \geq 1} \Gamma_s^{\mathbf{y}}(M) \oplus \Gamma_{\mathbf{x}}^{t}(M),$$

where

$$\Gamma_s^{\mathbf{y}}(M) = \begin{cases} \mathbf{M}_1^{\mathbf{y}}, & s = 1;\ \mathbf{y} = \mathbf{y}_q^{n,j} \text{ for some } n, j, q \\ \left\langle A \in \mathbf{M}_s^{\mathbf{y}} \mid \hat{e}_A \subset \Delta^{(q-1)}(P_{s-1}) \right\rangle, & s \geq 2 \\ 0, & \text{otherwise,} \end{cases}$$

$$\Gamma_{\mathbf{x}}^{t}(M) = \begin{cases} \mathbf{M}_{\mathbf{x}}^{1}, & t = 1;\ \mathbf{x} = \mathbf{x}_{m,i}^{p} \text{ for some } m, i, p \\ \left\langle B \in \mathbf{M}_{\mathbf{x}}^{t} \mid \check{e}_B \subset \Delta^{(p-1)}(P_{t-1}) \right\rangle, & t \geq 2 \\ 0, & \text{otherwise.} \end{cases}$$

Thus $\Gamma_{\mathbf{x}}^{t}(M)$ is generated by those tensor monomials $B = \beta_{x_1}^{t} \otimes \cdots \otimes \beta_{x_p}^{t} \in M_{t,x_1} \otimes \cdots \otimes M_{t,x_p}$ whose tensor factor $\beta_{x_i}^{t}$ is identified with some factor of a product cell in $\Delta^{(p-1)}(P_{t-1})$ corresponding to some column factorization $\beta_{x_i}^{t} = B_{i,l} \cdots B_{i,1}$ with respect to $\mathbf{W}$, and dually for $\Gamma_s^{\mathbf{y}}(M)$.

Definition 8. *A local prematrad* $(M, \gamma_{\mathbf{W}}, \eta)$ *is a **(left) matrad** if*

$$\Gamma^{\mathbf{y}}_{p}(M, \gamma_{\mathbf{W}}) \otimes \Gamma^{q}_{\mathbf{x}}(M, \gamma_{\mathbf{W}}) = \mathbf{W}^{\mathbf{y}}_{p} \otimes \mathbf{W}^{q}_{\mathbf{x}}$$

for all $p, q \geq 2$. *A **morphism of matrads** is a map of underlying local prematrads.*

The domain of the free prematrad $(M = F^{\text{pre}}(\Theta), \gamma,\ \eta)$ generated by $\Theta = \langle \theta^n_m \rangle_{m,n\geq 1}$ is $\mathbf{V} = M \oplus \bigoplus_{\mathbf{x},\mathbf{y}\notin \mathbb{N};\ s,t\in\mathbb{N}} \mathbf{M}^{\mathbf{y}}_{s} \oplus \mathbf{M}^{t}_{\mathbf{x}}$, whose submodules M, $\mathbf{M}^{1}_{\mathbf{x}}$, and $\mathbf{M}^{\mathbf{y}}_{1}$ are contained in the configuration module $\Gamma(M)$. As above, the symbol "$\cdot$" denotes the γ product.

Definition 9. *Let* $(M = F^{\text{pre}}(\Theta), \gamma, \eta)$ *be the free prematrad generated by* $\Theta = \langle \theta^n_m \rangle_{m,n\geq 1}$, *let* $F(\Theta) = \Gamma(M)\cdot\Gamma(M)$, *and let* $\gamma_{F(\Theta)} = \gamma|_{\Gamma(M)\otimes\Gamma(M)}$. *The **free matrad generated by** Θ is the triple* $\left(F(\Theta), \gamma_{F(\Theta)}, \eta\right)$.

The basis elements of $F_{n,m}(\Theta)$ lie in one-to-one correspondence with the cells of the polytope $KK_{n,m}$ mentioned in the introduction, and there is a differential ∂ of degree -1 on $F_{n,m}(\Theta)$ that agrees with the cellular differential on the cellular chain complex $C_*(KK_{n,m})$. We denote the differential object $(F(\Theta), \partial)$ by $\mathcal{H}_\infty$, and define an A_∞-infinity bialgebra as an algebra over the matrad $\mathcal{H}_\infty$.

3. RELATIVE MATRADS

Let R be a commutative ring (graded or ungraded) with unity 1_R, let (M, γ_M, η_M) and (N, γ_N, η_N) be R-prematrads, and let $E = \{E_{n,m}\}_{m,n\geq 1}$ be a bigraded R-module. Left and right actions $\lambda = \{\lambda^{\mathbf{y}}_{\mathbf{x}} : \mathbf{M}^{\mathbf{y}}_{p} \otimes \mathbf{E}^{q}_{\mathbf{x}} \to \mathbf{E}^{|\mathbf{y}|}_{|\mathbf{x}|}\}$ and $\rho = \{\rho^{\mathbf{y}}_{\mathbf{x}} : \mathbf{E}^{\mathbf{y}}_{p} \otimes \mathbf{N}^{q}_{\mathbf{x}} \to \mathbf{E}^{|\mathbf{y}|}_{|\mathbf{x}|}\}$ induce left and right upsilon products $\Upsilon_\lambda : \overline{\mathbf{M}} \otimes \overline{\mathbf{E}} \to \overline{\mathbf{E}}$ and $\Upsilon_\rho : \overline{\mathbf{E}} \otimes \overline{\mathbf{N}} \to \overline{\mathbf{E}}$ in the same way that $\gamma_M = \{\gamma^{\mathbf{y}}_{\mathbf{x}} : \mathbf{M}^{\mathbf{y}}_{p} \otimes \mathbf{M}^{q}_{\mathbf{x}} \to \mathbf{M}^{|\mathbf{y}|}_{|\mathbf{x}|}\}$ induces an upsilon product $\Upsilon_M : \overline{\mathbf{M}} \otimes \overline{\mathbf{M}} \to \overline{\mathbf{M}}$ (see [22] and formula (2.1)).

Definition 10. *A tuple* (M, E, N, λ, ρ) *is a **relative prematrad** if*

- *(i) Associativity holds:*
  - *(a)* $\Upsilon_\rho(\Upsilon_\lambda \otimes \mathbf{1}) = \Upsilon_\lambda(\mathbf{1} \otimes \Upsilon_\rho)$;
  - *(b)* $\Upsilon_\lambda(\Upsilon_M \otimes \mathbf{1}) = \Upsilon_\lambda(\mathbf{1} \otimes \Upsilon_\lambda)$;
  - *(c)* $\Upsilon_\rho(\Upsilon_\rho \otimes \mathbf{1}) = \Upsilon_\rho(\mathbf{1} \otimes \Upsilon_N)$.
- *(ii) The units* η_M *and* η_N *induce the following canonical isomorphisms for all* $a, b \in \mathbb{N}$*:*

$$R^{\otimes b} \otimes \mathbf{E}^{b}_{a} \xrightarrow{\eta^{\otimes b}_{M} \otimes \mathbf{1}} \mathbf{M}^{\mathbf{1}^{b\times 1}}_{1} \otimes \mathbf{E}^{b}_{a} \xrightarrow{\lambda^{\mathbf{1}^{b\times 1}}_{a}} \mathbf{E}^{b}_{a};$$

$$\mathbf{E}^{b}_{a} \otimes R^{\otimes a} \xrightarrow{\mathbf{1} \otimes \eta^{\otimes a}_{N}} \mathbf{E}^{b}_{a} \otimes \mathbf{N}^{1}_{\mathbf{1}^{1\times a}} \xrightarrow{\rho^{b}_{\mathbf{1}^{1\times a}}} \mathbf{E}^{b}_{a}.$$

We refer to E *as an* M-N-***bimodule;*** *when* $M = N$ *we refer to* E *as an* M-***bimodule***.

Definition 11. *A **morphism*** $f : (M, E, N, \lambda, \rho) \to (M', E', N', \lambda', \rho')$ *of relative prematrads is a triple* $(f_M : M \to M',\ f_E : E \to E',\ f_N : N \to N')$ *such that*

- *(i)* f_M *and* f_N *are maps of prematrads;*
- *(ii)* f_E *commutes with left and right actions, i.e.,* $f_E \circ \lambda^{\mathbf{y}}_{\mathbf{x}} = \lambda'^{\mathbf{y}}_{\mathbf{x}} \circ (f^{\otimes q}_{M} \otimes f^{\otimes p}_{E})$ *and* $f_E \circ \rho^{\mathbf{y}}_{\mathbf{x}} = \rho'^{\mathbf{y}}_{\mathbf{x}} \circ (f^{\otimes q}_{E} \otimes f^{\otimes p}_{N})$ *for all* $\mathbf{x} \times \mathbf{y} \in \mathbb{N}^{p\times 1} \times \mathbb{N}^{1\times q}$.

Tree representations of $\lambda_1^{\mathbf{x}}$ and $\rho_1^{\mathbf{x}}$ are related to those of $\lambda_{\mathbf{x}}^1$ and $\rho_{\mathbf{x}}^1$ by a reflection in some horizontal axis. Although $\rho_{\mathbf{x}}^1$ agrees with Markl, Shnider, and Stasheff's right module action over an operad [16], $\lambda_{\mathbf{x}}^1$ differs fundamentally from their left module action, and our definition of an "operadic bimodule" is consistent with their definition of an operadic ideal.

Given graded R-modules A and B, let

$$\begin{array}{lllll} U_A & = & \mathcal{E}nd_{TA} & = & \{N_{s,p} = Hom\,(A^{\otimes p}, A^{\otimes s})\}_{p,s\geq 1} \\ U_{A,B} & = & Hom\,(TA, TB) & = & \{E_{t,q} = Hom\,(A^{\otimes q}, B^{\otimes t})\}_{q,t\geq 1} \\ U_B & = & \mathcal{E}nd_{TB} & = & \{M_{u,r} = Hom\,(B^{\otimes r}, B^{\otimes u})\}_{r,u\geq 1}\,, \end{array}$$

and define left and right actions λ and ρ in terms of the horizontal and vertical operations $\times$ and $\circ$ analogous to those in the prematrad structures on U_A and U_B (see Section 2 above and [22]). Then the relative PROP $(U_B, U_{A,B}, U_A, \lambda, \rho)$ is the universal example of a relative prematrad.

We adopt the notation in Definition 4: $\overline{\mathbf{G}}$ denotes the set of matrices over $G^{\mathrm{pre}} = G^{\mathrm{pre}}_{*,*}$ and $\mathbf{G}$ denotes the subset of bisequence matrices in $\overline{\mathbf{G}}$. Given sets $\mathcal{G}^{\mathrm{pre}}_{j,i}$, $i \leq m$, $j \leq n$, $i + j < m + n$, let $\overline{\mathbf{C}}_{[n,m]}$ denote the set of matrices over

$$\mathcal{G}_{[n,m]} = \bigcup_{\substack{i\leq m,\ j\leq n,\\ i+j<m+n}} \mathcal{G}^{\mathrm{pre}}_{j,i}.$$

If $\mathcal{G}^{\mathrm{pre}}_{n,m}$ is given for each $m, n \geq 1$, then $\overline{\mathbf{C}}$ denotes the set of matrices over $\mathcal{G}^{\mathrm{pre}} = \mathcal{G}^{\mathrm{pre}}_{*,*}$ and $\mathbf{C}$ denotes the subset of bisequence matrices in $\overline{\mathbf{C}}$.

Definition 12. *Given a free bigraded R-module $\Theta = \left\langle \theta^n_m \mid \theta^1_1 = \mathbf{1} \right\rangle_{m,n\geq 1}$ generated by singletons in each bidegree, let $(F^{\mathrm{pre}}(\Theta), \gamma, \eta)$ be the free prematrad generated by Θ. Let $\mathfrak{F} = \langle \mathfrak{f}^n_m \rangle_{m,n\geq 1}$ be a free bigraded R-module generated by singletons in each bidegree and set $\mathcal{G}^{\mathrm{pre}}_{1,1} = \mathfrak{f}^1_1$. Inductively, if $m + n \geq 3$ and $\overline{\mathbf{C}}_{[n,m]}$ has been constructed, define*

$$\mathcal{G}^{\mathrm{pre}}_{n,m} = \mathfrak{f}^n_m \cup \{B_1 \cdots B_l \cdot C \cdot A_k \cdots A_1 \mid k + l \geq 1;\ k, l \geq 0\}\,,$$

where

(i) $A_1 \in \mathbf{G}^q_{\mathbf{x}}$ and $B_1 \in \mathbf{G}^{\mathbf{y}}_p$ for $(|\mathbf{x}|, |\mathbf{y}|) = (m, n)$;

(ii) $A_i, B_j \in \overline{\mathbf{G}}_{[n,m]}$ for all i, j;

(iii) $C \in \overline{\mathbf{C}}_{[n,m]}$; and

(vii) some association of $B_1 \cdots B_l \cdot C \cdot A_k \cdots A_1$ defines a sequence of BTPs.

Let $\sim$ be the equivalence relation on $\mathcal{G}^{\mathrm{pre}} = \mathcal{G}^{\mathrm{pre}}_{*,*}$ generated by

$$[X_{ij}Y_{ij}] \sim [X_{ij}]\,[Y_{ij}] \text{ iff } [X_{ij}] \times [Y_{ij}] \in \overline{\mathbf{C}} \times \overline{\mathbf{G}} \cup \overline{\mathbf{G}} \times \overline{\mathbf{G}} \cup \overline{\mathbf{G}} \times \overline{\mathbf{C}} \text{ is a BTP,}$$

and let $F^{\mathrm{pre}}(\Theta, \mathfrak{F}, \Theta) = \langle \mathcal{G}^{\mathrm{pre}} / \sim \rangle$. The ***free relative prematrad generated by*** *Θ* ***and*** *$\mathfrak{F}$ is the relative prematrad*

$$(F^{\mathrm{pre}}(\Theta), F^{\mathrm{pre}}(\Theta, \mathfrak{F}, \Theta), F^{\mathrm{pre}}(\Theta), \lambda^{\mathrm{pre}}, \rho^{\mathrm{pre}}),$$

where λ^{pre} and ρ^{pre} are juxtaposition.

Example 1. *The module $F^{\mathrm{pre}}_{2,2}(\Theta, \mathfrak{F}, \Theta)$ contains 25 module generators, namely, the indecomposable $\mathfrak{f}^2_2$ and the following $(\lambda^{\mathrm{pre}}, \rho^{\mathrm{pre}})$-decomposables:*

Two of the form BCA:

$$[\theta_1^2]\,[\mathfrak{f}_1^1]\,[\theta_2^1] \;\; \text{and} \; \begin{bmatrix} \theta_2^1 \\ \theta_2^1 \end{bmatrix} \begin{bmatrix} \mathfrak{f}_1^1 & \mathfrak{f}_1^1 \\ \mathfrak{f}_1^1 & \mathfrak{f}_1^1 \end{bmatrix} \begin{bmatrix} \theta_1^2 & \theta_1^2 \end{bmatrix}. \tag{3.1}$$

Eleven of the form $CA_k \cdots A_1$:

$$[\mathfrak{f}_1^2]\,[\theta_2^1], \; \begin{bmatrix} \mathfrak{f}_1^1 \\ \mathfrak{f}_1^1 \end{bmatrix} [\theta_2^2], \; \begin{bmatrix} \mathfrak{f}_1^1 \\ \mathfrak{f}_1^1 \end{bmatrix} [\theta_1^2]\,[\theta_2^1], \; \begin{bmatrix} \mathfrak{f}_1^1 \\ \mathfrak{f}_1^1 \end{bmatrix} \begin{bmatrix} \theta_2^1 \\ \theta_2^1 \end{bmatrix} \begin{bmatrix} \theta_1^2 & \theta_1^2 \end{bmatrix},$$

$$\begin{bmatrix} \mathfrak{f}_2^1 \\ \mathfrak{f}_2^1 \end{bmatrix} \begin{bmatrix} \theta_1^2 & \theta_1^2 \end{bmatrix}, \; \begin{bmatrix} \mathfrak{f}_2^1 \\ [\mathfrak{f}_1^1]\,[\theta_2^1] \end{bmatrix} \begin{bmatrix} \theta_1^2 & \theta_1^2 \end{bmatrix}, \; \begin{bmatrix} [\mathfrak{f}_1^1]\,[\theta_2^1] \\ \mathfrak{f}_2^1 \end{bmatrix} \begin{bmatrix} \theta_1^2 & \theta_1^2 \end{bmatrix},$$

$$\begin{bmatrix} [\theta_2^1] \begin{bmatrix} \mathfrak{f}_1^1 & \mathfrak{f}_1^1 \end{bmatrix} \\ [\mathfrak{f}_1^1]\,[\theta_2^1] \end{bmatrix} \begin{bmatrix} \theta_1^2 & \theta_1^2 \end{bmatrix}, \; \begin{bmatrix} [\mathfrak{f}_1^1]\,[\theta_2^1] \\ [\theta_2^1] \begin{bmatrix} \mathfrak{f}_1^1 & \mathfrak{f}_1^1 \end{bmatrix} \end{bmatrix} \begin{bmatrix} \theta_1^2 & \theta_1^2 \end{bmatrix},$$

$$\begin{bmatrix} [\theta_2^1] \begin{bmatrix} \mathfrak{f}_1^1 & \mathfrak{f}_1^1 \end{bmatrix} \\ \mathfrak{f}_2^1 \end{bmatrix} \begin{bmatrix} \theta_1^2 & \theta_1^2 \end{bmatrix}, \; \begin{bmatrix} \mathfrak{f}_2^1 \\ [\theta_2^1] \begin{bmatrix} \mathfrak{f}_1^1 & \mathfrak{f}_1^1 \end{bmatrix} \end{bmatrix} \begin{bmatrix} \theta_1^2 & \theta_1^2 \end{bmatrix}.$$

And eleven respective duals of the form $B_1 \cdots B_k C$.

Example 2. *Recall that the bialgebra prematrad $\mathcal{H}^{\mathrm{pre}}$ has two prematrad generators $c_{1,2}$ and $c_{2,1}$, and a single module generator $c_{n,m}$ in bidegree (m,n) (see* [22]*). Consequently, the $\mathcal{H}^{\mathrm{pre}}$-bimodule $\mathcal{JJ}^{\mathrm{pre}}$ has a single bimodule generator $\mathfrak{f}$ of bidegree $(1,1)$ and a single module generator in each bidegree satisfying the structure relations*

$$\lambda(c_{n,m}; \underbrace{\mathfrak{f}, \ldots, \mathfrak{f}}_{m}) = \rho(\underbrace{\mathfrak{f}, \ldots, \mathfrak{f}}_{n}; c_{n,m}).$$

More precisely, if $\Theta = \langle \theta_1^1 = \mathbf{1}, \theta_2^1, \theta_1^2 \rangle$ and $\mathfrak{F} = \langle \mathfrak{f} = \mathfrak{f}_1^1 \rangle$, then

$$\mathcal{JJ}^{\mathrm{pre}} = F^{\mathrm{pre}}(\Theta, \mathfrak{F}, \Theta) \,/ \sim$$

with $A \sim B$ if and only if $\mathrm{bideg}(A) = \mathrm{bideg}(B)$ for $A, B \in F^{\mathrm{pre}}(\Theta, \mathfrak{F}, \Theta)$.

A morphism $f : A \to B$ of bialgebras is the image of $\mathfrak{f}$ under a map $\mathcal{JJ}^{\mathrm{pre}} \to U_{A,B}$ of relative prematrads.

Example 3. *Whereas $F_{1,*}^{\mathrm{pre}}(\Theta)$ and $F_{*,1}^{\mathrm{pre}}(\Theta)$ can be identified with the A_∞-operad $\mathcal{A}_\infty$ (see* [22]*), $F_{1,*}(\Theta, \mathfrak{F}, \Theta)$ and $F_{*,1}(\Theta, \mathfrak{F}, \Theta)$ can be identified with the $\mathcal{A}_\infty$-bimodule $\mathcal{J}_\infty$ whose bimodule generators are in 1-1 correspondence with $\{\mathfrak{f}_m^1\}_{m \geq 1}$ and $\{\mathfrak{f}_1^n\}_{n \geq 1}$, respectively. Thus a morphism $f : A \to B$ of A_∞-(co)algebras is the image of $\mathcal{A}_\infty$-bimodule generators under a map of relative prematrads $\mathcal{J}_\infty \to Hom\,(TA, B)$ (or $\mathcal{J}_\infty \to Hom\,(A, TB)$).*

When $\Theta = \left\langle \theta_m^n \neq 0 \mid \theta_1^1 = \mathbf{1} \right\rangle_{m,n \geq 1}$ and $\mathfrak{F} = \langle \mathfrak{f}_m^n \neq 0 \rangle_{m,n \geq 1}$, the canonical projections $\varrho_\Theta^{\mathrm{pre}} : F^{\mathrm{pre}}(\Theta) \to \mathcal{H}^{\mathrm{pre}}$ and $\varrho^{\mathrm{pre}} : F^{\mathrm{pre}}(\Theta, \mathfrak{F}, \Theta) \to \mathcal{JJ}^{\mathrm{pre}}$ give a map $(\varrho_\Theta^{\mathrm{pre}}, \varrho^{\mathrm{pre}}, \varrho_\Theta^{\mathrm{pre}})$ of relative prematrads. If ∂^{pre} is a differential on $F^{\mathrm{pre}}(\Theta, \mathfrak{F}, \Theta)$

such that ϱ^{pre} is a free resolution in the category of relative prematrads, the induced isomorphism $\varrho^{\text{pre}} : H_*\left(F^{\text{pre}}(\Theta,\mathfrak{F},\Theta),\partial^{\text{pre}}\right) \approx \mathcal{J}\mathcal{J}^{\text{pre}}$ implies

$$\partial^{\text{pre}}(\mathfrak{f}_1^1) = 0$$

$$\partial^{\text{pre}}(\mathfrak{f}_2^1) = \rho(\mathfrak{f}_1^1;\theta_2^1) - \lambda(\theta_2^1;\mathfrak{f}_1^1,\mathfrak{f}_1^1)$$

$$\partial^{\text{pre}}(\mathfrak{f}_1^2) = \rho(\mathfrak{f}_1^1,\mathfrak{f}_1^1;\theta_1^2) - \lambda(\theta_1^2;\mathfrak{f}_1^1).$$

This gives rise to the standard isomorphisms

$$\begin{array}{ccccccc}
F_{1,2}^{\text{pre}}(\Theta,\mathfrak{F},\Theta) & = & \big\langle \left[\theta_2^1\right]\left[\begin{array}{cc}\mathfrak{f}_1^1 & \mathfrak{f}_1^1\end{array}\right] & , & \left[\mathfrak{f}_2^1\right] & , & \left[\mathfrak{f}_1^1\right]\left[\theta_2^1\right]\big\rangle \\
\approx\ \updownarrow & & \updownarrow & & \updownarrow & & \updownarrow \\
C_*(J_2) & = & \langle 1|2 & , & 12 & , & 2|1\rangle \\
\approx\ \updownarrow & & \updownarrow & & \updownarrow & & \updownarrow \\
F_{2,1}^{\text{pre}}(\Theta,\mathfrak{F},\Theta) & = & \big\langle \left[\theta_1^2\right]\left[\mathfrak{f}_1^1\right] & , & \left[\mathfrak{f}_1^2\right] & , & \left[\begin{array}{c}\mathfrak{f}_1^1 \\ \mathfrak{f}_1^1\end{array}\right]\left[\theta_1^2\right]\big\rangle
\end{array}$$

(see Figures 3 and 4). A similar application of ∂^{pre} to $\mathfrak{f}_1^n$ and $\mathfrak{f}_m^1$ gives the isomorphisms

$$F_{n,1}^{\text{pre}}(\Theta,\mathfrak{F},\Theta) \xrightarrow{\approx} \mathcal{J}_\infty(n) = C_*(J_n) \tag{3.2}$$

and

$$F_{1,m}^{\text{pre}}(\Theta,\mathfrak{F},\Theta) \xrightarrow{\approx} \mathcal{J}_\infty(m) = C_*(J_m) \tag{3.3}$$

(see [23], [24], [20]). As in the absolute case, there is a differential ∂ on a canonical proper submodule $\mathcal{J}\mathcal{J}_\infty \subset F^{\text{pre}}(\Theta,\mathfrak{F},\Theta)$ such that the canonical projection $\varrho : \mathcal{J}\mathcal{J}_\infty \to \mathcal{J}\mathcal{J}^{\text{pre}}$ is a free resolution in the category of "relative matrads."

Definition 13. *Given local prematrads* $(M,\gamma_{\mathbf{W}_M},\eta_M)$ *and* $(N,\gamma_{\mathbf{W}_N},\eta_N)$, *a bigraded R-module* $E=\{E_{n,m}\}_{m,n\geq 1}$, *a telescoping submodule* $\mathbf{W}_E \subseteq \mathbf{V}_E$, *and actions* $\lambda=\{\lambda_{\mathbf{x}}^{\mathbf{y}}:(\mathbf{W}_M)_p^{\mathbf{y}}\otimes(\mathbf{W}_E)_{\mathbf{x}}^q \to (\mathbf{W}_E)_{|\mathbf{x}|}^{|\mathbf{y}|}\}$ *and* $\rho=\{\rho_{\mathbf{x}}^{\mathbf{y}}:(\mathbf{W}_E)_p^{\mathbf{y}}\otimes(\mathbf{W}_N)_{\mathbf{x}}^q \to (\mathbf{W}_E)_{|\mathbf{x}|}^{|\mathbf{y}|}\}$, *the tuple* (M,E,N,λ,ρ) *is a* ***relative local prematrad with domain*** $(\mathbf{W}_M,\mathbf{W}_E,\ \mathbf{W}_N)$ *if*

(i) $(\mathbf{W}_E)_{\mathbf{x}}^1=\mathbf{E}_{\mathbf{x}}^1$ *and* $(\mathbf{W}_E)_1^{\mathbf{y}}=\mathbf{E}_1^{\mathbf{y}}$ *for all* $\mathbf{x},\mathbf{y}$;
(ii) $\Upsilon_M,\Upsilon_N,\Upsilon_\lambda,\Upsilon_\rho$ *interact associatively on* $\mathcal{W}_E\cap\mathbf{E}$;
(iii) the relative prematrad unit axiom holds for λ *and* ρ.

Let

$$\overline{\mathbf{E}}_{\text{row}} = \bigoplus_{\mathbf{x},Y\in\mathbb{N}^{q\times p};\ p,q\geq 1} E_{Y,\mathbf{x}} \quad\text{and}\quad \overline{\mathbf{E}}^{\text{col}} = \bigoplus_{X,\mathbf{y}\in\mathbb{N}^{q\times p};\ p,q\geq 1} E_{\mathbf{y},X},$$

where $\mathbf{x}$ has constant columns and $\mathbf{y}$ has constant rows. If A is a $q\times p$ monomial in $E_{Y,\mathbf{x}}$, we refer to a row of $\mathbf{x}$ as the *row leaf sequence* of A and write $\mathbf{x}=$rls$A$; if B is a $q\times p$ monomial in $E_{\mathbf{y},X}$, we refer to a column of $\mathbf{y}$ as the *column leaf sequence* of B and write $\mathbf{y}=$cls$B$. Given a telescoping submodule $\mathbf{W}\subseteq\mathbf{V}_E$ and its telescopic extension $\mathcal{W}_E$, let $(\mathcal{W}_E)_{\text{row}}=\mathcal{W}_E\cap\overline{\mathbf{E}}_{\text{row}}$ and $(\mathcal{W}_E)^{\text{col}}=\mathcal{W}_E\cap\overline{\mathbf{E}}^{\text{col}}$.

Definition 14. *Let* (M, E, N, λ, ρ) *be a relative local prematrad with domain* $(\mathbf{W}_M, \mathbf{W}_E, \mathbf{W}_N)$, *let* $m, n \geq 1$, *let* $\mathfrak{f} \in E_{*,m}$ *and* $\mathfrak{g} \in E_{n,*}$.

(i) *A* ***row factorization of*** $\mathfrak{f}$ ***with respect to*** $(\mathbf{W}_M, \mathbf{W}_E, \mathbf{W}_N)$ *is an* $(\Upsilon_\lambda, \Upsilon_\rho)$*-factorization*

$$A_1 \cdots A_s \cdot C \cdot A'_{s+1} \cdots A'_k = \mathfrak{f}$$

such that

(a) $A_i \in (\mathcal{W}_M)_{\text{row}}$ *and* $\text{rls}A_i \neq \mathbf{1}$ *for all* i;

(b) $C \in (\mathcal{W}_E)_{\text{row}}$;

(c) $A'_j \in (\mathcal{W}_N)_{\text{row}}$ *and* $\text{rls}A'_j \neq \mathbf{1}$ *for all* j.

The $(k+1)$*-tuple of row vectors*

$$\mathbf{m} = \left(\text{rls}A_1, \ldots, \text{rls}A_s, (1, 0, \ldots, 0) + \text{rls}C, \left(1, \text{rls}A'_{s+1}\right), \ldots, \left(1, \text{rls}A'_k\right)\right)$$

is the related $(m+1, k+1)$*-**row descent sequence** of* $\mathfrak{f}$.

(ii) *A* ***column factorization of*** $\mathfrak{g}$ ***with respect to*** $(\mathbf{W}_M, \mathbf{W}_E, \mathbf{W}_N)$ *is an* $(\Upsilon_\lambda, \Upsilon_\rho)$*-factorization*

$$B'_l \cdots B'_{t+1} \cdot D \cdot B_t \cdots B_1 = \mathfrak{g}$$

such that

(a) $B'_j \in (\mathcal{W}_M)^{\text{col}}$ *and* $\text{cls}B'_j \neq \mathbf{1}$ *for all* j;

(b) $D \in (\mathcal{W}_E)^{\text{col}}$;

(c) $B_j \in (\mathcal{W}_N)^{\text{col}}$ *and* $\text{cls}B_j \neq \mathbf{1}$ *for all* j.

The $(l+1)$*-tuple of column vectors*

$$\mathbf{n} = \left(\begin{pmatrix} \text{cls}B'_l \\ 1 \end{pmatrix}, \ldots, \begin{pmatrix} \text{cls}B'_{t+1} \\ 1 \end{pmatrix}, \text{cls}D + \begin{pmatrix} 0 \\ \vdots \\ 0 \\ 1 \end{pmatrix}, \text{cls}B_t, \ldots, \text{cls}B_1 \right)$$

is the related $(n+1, l+1)$*-**column descent sequence of*** $\mathfrak{g}$.

Column and row factorizations are not unique. Note that $\mathfrak{f} \in E_{n,m}$ always has trivial row and column factorizations as a 1×1 matrix $C = [\mathfrak{f}]$.

Given elements $\mathfrak{f} \in E_{*,m}$ and $\mathfrak{g} \in E_{n,*}$ with $m, n \geq 1$, choose a row factorization $A_1 \cdots A_s \cdot C \cdot A'_{s+1} \cdots A'_k$ of $\mathfrak{f}$ and a column factorization $B'_l \cdots B'_{t+1} \cdot D \cdot B_t \cdots B_1$ of $\mathfrak{g}$. The related row descent sequence $\mathbf{m}$ identifies $\mathfrak{f}$ with an up-rooted $(m+1)$-leaf, $(k+1)$-level planar rooted tree with levels (PLT) and hence with a codimension k face $\hat{e}_{\mathfrak{f}}$ of P_m. Dually, the related column descent sequence $\mathbf{n}$ identifies $\mathfrak{g}$ with a down-rooted $(n+1)$-leaf, $(l+1)$-level PLT and hence with a codimension l face $\overset{\vee}{e}_{\mathfrak{g}}$ of P_n. Extending to Cartesian products, identify the monomials $F = \mathfrak{f}_1 \otimes \cdots \otimes \mathfrak{f}_q \in (E_{*,m})^{\otimes q}$ and $G = \mathfrak{g}_1 \otimes \cdots \otimes \mathfrak{g}_p \in (E_{n,*})^{\otimes p}$ with the product cells

$$\hat{e}_F = \hat{e}_{\mathfrak{f}_1} \times \cdots \times \hat{e}_{\mathfrak{f}_q} \subset P_m^{\times q} \quad \text{and} \quad \overset{\vee}{e}_G = \overset{\vee}{e}_{\mathfrak{g}_1} \times \cdots \times \overset{\vee}{e}_{\mathfrak{g}_p} \subset P_n^{\times p}.$$

As in the absolute case discussed in Section 2, the product cell $\hat{e}_F$ either *is* or *is not* a subcomplex of $\Delta^{(q-1)}(P_m) \subset P_m^{\times q}$, and dually for $\overset{\vee}{e}_G$. Recall that $\mathbf{x} \times \mathbf{y} \in \mathbb{N}^{1 \times p} \times \mathbb{N}^{q \times 1}$ and let

$$r\Gamma(E) = E \oplus \bigoplus_{\mathbf{x}, \mathbf{y} \notin \mathbb{N};\ s,t \geq 1} r\Gamma_s^{\mathbf{y}}(E) \oplus r\Gamma_{\mathbf{x}}^t(E),$$

where

$$r\Gamma_s^{\mathbf{y}}(E) = \left\langle F \in \mathbf{E}_s^{\mathbf{y}} \mid \hat{e}_F \subset \Delta^{(q-1)}\left(P_s\right)\right\rangle,$$

$$r\Gamma_{\mathbf{x}}^t(E) = \left\langle G \in \mathbf{E}_{\mathbf{x}}^t \mid \check{e}_G \subset \Delta^{(p-1)}\left(P_t\right)\right\rangle.$$

Definition 15. *The* ***(left) configuration module*** *of a relative local prematrad (M, E, N, λ, ρ) with domain $(\mathbf{W}_M, \mathbf{W}_E, \mathbf{W}_N)$ is the triple*

$$\left(\Gamma(M, \gamma_{\mathbf{W}_M}),\ r\Gamma(E),\ \Gamma(N, \gamma_{\mathbf{W}_N})\right).$$

When $s = t = 1$, the arguments in the absolute case carry over verbatim and give

$$\bigoplus_{\mathbf{x}} r\Gamma_{\mathbf{x}}^1(E) = T^+(E_{1,*}) \quad \text{and} \quad \bigoplus_{\mathbf{y}} r\Gamma_1^{\mathbf{y}}(E) = T^+(E_{*,1}).$$

Definition 16. *Let $(M, \gamma_{\mathbf{W}_M})$ and $(N, \gamma_{\mathbf{W}_N})$ be (left) matrads. A relative local prematrad (M, E, N, λ, ρ) with domain $(\mathbf{W}_M, \mathbf{W}_E, \mathbf{W}_N)$ is a* ***relative (left) matrad*** *if $r\Gamma(E) = \mathbf{W}_E$. When $M = N$ we refer to $r\Gamma(E)$ as a* ***$\Gamma(M)$-bimodule****. A* ***morphism of relative matrads*** *is a map of underlying relative local prematrads.*

A relative prematrad (M, E, N, ρ, λ) with domain $(\mathbf{W}_M, \mathbf{W}_E, \mathbf{W}_N)$ restricts to a relative matrad structure with domain $(\Gamma(M), r\Gamma(E), \Gamma(N))$.

Example 4. The Bialgebra Morphism Matrad $\mathcal{JJ}$. *The $\mathcal{H}^{\text{pre}}$-bimodule $\mathcal{JJ}^{\text{pre}}$ discussed in Example 2 satisfies*

$$r\Gamma_p^{\mathbf{y}}(E) \otimes \Gamma_{\mathbf{x}}^q(M) = \mathbf{E}_p^{\mathbf{y}} \otimes \mathbf{M}_{\mathbf{x}}^q \quad \text{and} \quad \Gamma_s^{\mathbf{v}}(M) \otimes r\Gamma_{\mathbf{u}}^t(E) = \mathbf{M}_s^{\mathbf{v}} \otimes \mathbf{E}_{\mathbf{u}}^t.$$

Consequently, $\mathcal{JJ}^{\text{pre}}$ is also a relative matrad, called ***the bialgebra morphism matrad****, and is henceforth denoted by $\mathcal{JJ}$.*

Definition 17. *Let $\Theta = \left\langle \theta_m^n \mid \theta_1^1 = \mathbf{1}\right\rangle_{m,n\geq 1}$ and $\mathfrak{F} = \left\langle \mathfrak{f}_m^n \right\rangle_{m,n\geq 1}$. Let $F^{\text{pre}} = F^{\text{pre}}(\Theta)$ and consider the free F^{pre}-bimodule $E^{\text{pre}} = F^{\text{pre}}\left(\Theta, \mathfrak{F}, \Theta\right)$. The* ***free relative matrad generated by $(\Theta, \mathfrak{F})$ with domain*** *$(\mathbf{W}_{F(\Theta)}, \mathbf{W}_E, \mathbf{W}_{F(\Theta)})$ is the tuple $\left(F\left(\Theta\right), F\left(\Theta, \mathfrak{F}, \Theta\right), F\left(\Theta\right), \lambda, \rho\right)$, where*

$$F\left(\Theta, \mathfrak{F}, \Theta\right) = \lambda^{\text{pre}}\left[\Gamma_p^{\mathbf{y}}(F^{\text{pre}}); r\Gamma_{\mathbf{x}}^q(E^{\text{pre}})\right] \oplus \rho^{\text{pre}}\left[r\Gamma_p^{\mathbf{y}}(E^{\text{pre}}); \Gamma_{\mathbf{x}}^q(F^{\text{pre}})\right],$$

$$\lambda = \lambda^{\text{pre}}|_{\mathbf{W}_{F(\Theta)} \otimes \mathbf{W}_E}, \quad \rho = \rho^{\text{pre}}|_{\mathbf{W}_E \otimes \mathbf{W}_{F(\Theta)}}, \quad \textit{and}$$

$$\mathbf{W}_E = \bigoplus_{p,q\in\mathbb{N};\ \mathbf{x},\mathbf{y}\notin\mathbb{N}} F_{q,p}(\Theta, \mathfrak{F}, \Theta) \oplus \Gamma_p^{\mathbf{y}}(E^{\text{pre}}) \oplus \Gamma_{\mathbf{x}}^q(E^{\text{pre}}).$$

For example, the monomials in (3.1) are two of the 17 module generators in $F_{2,2}\left(\Theta, \mathfrak{F}, \Theta\right)$ identified with the faces of the octagon $JJ_{2,2}$ (see Figure 5). In general, $F_{1,m}\left(\Theta, \mathfrak{F}, \Theta\right) = F_{1,m}^{\text{pre}}\left(\Theta, \mathfrak{F}, \Theta\right)$ and $F_{n,1}\left(\Theta, \mathfrak{F}, \Theta\right) = F_{n,1}^{\text{pre}}\left(\Theta, \mathfrak{F}, \Theta\right)$ for all $m, n \geq 1$.

Example 5. The A_∞-bialgebra Morphism Matrad**.** *Let $\Theta = \left\langle \theta_m^n \neq 0 | \theta_1^1 = \mathbf{1}\right\rangle_{m,n\geq 1}$ and $\mathfrak{F} = \left\langle \mathfrak{f}_m^n \neq 0\right\rangle_{m,n\geq 1}$, where $|\mathfrak{f}_m^n| = m + n - 2$. Following the construction in the absolute case, let $r\mathcal{C}$ be the set indexing the module generators $F(\Theta, \mathfrak{F}, \Theta)$, and let $\mathcal{AR}_{m,n} \subset \mathcal{C}_{m,n} \times r\mathcal{C}_{m,n}$ and $\mathcal{QB}_{m,n} \subset r\mathcal{C}_{m,n} \times \mathcal{C}_{m,n}$ be the subsets of $r\mathcal{C}$ that index the codimension 1 elements of $F_{n,m}(\Theta, \mathfrak{F}, \Theta)$ of the form $\lambda(-;-)$ and $\rho(-;-)$, respectively. Let $\left\{(A_p^{\mathbf{y}})_\alpha\right\}$ and $\{(B_{\mathbf{x}}^q)_\beta\}$ be the bases defined in Example 19 of* [22]*, and let $\{(Q_s{}^{\mathbf{v}})_\nu\}_{\nu\in\mathcal{Q}_s^{\mathbf{v}}}$ and $\{(R_{\mathbf{u}}^t)_\mu\}_{\mu\in\mathcal{R}_{\mathbf{u}}^t}$ be the analogous bases for $r\Gamma_s^{\mathbf{v}}(E)$ and*

$r\Gamma^t_{\mathbf{u}}(E)$ in dimensions $|\mathbf{v}|+s-t$ and $|\mathbf{u}|+t-s$; then each $\hat{e}_{Q_\nu}$ is a subcomplex of $\Delta^{(t-1)}(P_s)$ with associated sign $(-1)^{\epsilon_\nu}$ and each $\overset{\vee}{e}_{R_\mu}$ is a subcomplex of $\Delta^{(s-1)}(P_t)$ with associated sign $(-1)^{\epsilon_\mu}$. Define a differential $\partial : F(\Theta,\mathfrak{F},\Theta)\to F(\Theta,\mathfrak{F},\Theta)$ of degree -1 on generators by

$$\begin{aligned}\partial(\mathfrak{f}^n_m) = &\sum_{(\alpha,\mu)\in\mathcal{AR}_{m,n}} (-1)^{\epsilon_1+\epsilon_\alpha+\epsilon_\mu}\lambda\left[(A^{\mathbf{y}}_p)_\alpha;(R^q_{\mathbf{x}})_\mu\right]\\ &+\sum_{(\nu,\beta)\in\mathcal{QB}_{m,n}} (-1)^{\epsilon_2+\epsilon_\nu+\epsilon_\beta}\rho\left[(Q^{\mathbf{y}}_p)_\nu;(B^q_{\mathbf{x}})_\beta\right],\end{aligned}\tag{3.4}$$

*where $(-1)^{\epsilon_1}$ is the sign associated with the codimension 1 cell $e_{(\mathbf{y},\mathbf{x})}=C|D\subset P_{m+n-2}$ defined in line (6.1) of [22], and $\epsilon_2=\#D+\epsilon_1+1$. Extend ∂ as a derivation of ρ and λ; then $\partial^2=0$ follows from the associativity of ρ and λ. The **$\mathcal{A}_\infty$-bialgebra morphism matrad is the** DG $\mathcal{H}_\infty$-bimodule $\mathcal{JJ}_\infty=(F(\Theta,\mathfrak{F},\Theta),\partial)$.*

We realize the $\mathcal{A}_\infty$-bialgebra morphism matrad by the cellular chains of a new family of polytopes $JJ=\sqcup_{m,n\geq1}JJ_{n,m}$, called *bimultiplihedra*, to be constructed in the next section. The standard isomorphisms (3.2) and (3.3) extend to isomorphisms

$$(\mathcal{JJ}_\infty)_{n,m}\xrightarrow{\approx}C_*(JJ_{n,m}),\tag{3.5}$$

and one recovers $\mathcal{J}_\infty$ by restricting the differential ∂ to $(\mathcal{JJ}_\infty)_{1,*}$ or $(\mathcal{JJ}_\infty)_{*,1}$.

4. Bimultiplihedra

In this section we construct the bimultiplihedron $JJ_{n,m}$ as a subdivision of the cylinder $KK_{n,m}\times I$. Whereas our construction of $KK_{n+1,m+1}$ uses the combinatorics of P_{m+n}, our construction of $JJ_{n+1,m+1}$ uses the combinatorics of P_{m+n+1} thought of as a subdivision of $P_{m+n}\times I$ (see [20]). And indeed, $JJ_{n+1,m+1}$ is the geometric realization of a poset $\mathcal{JJ}_{n+1,m+1}=r\mathcal{PP}_{n,m}/\sim$ whose construction resembles that of $\mathcal{KK}_{n+1,m+1}=\mathcal{PP}_{n,m}/\sim$ given in [22], but with some technical differences.

Recall that faces of the permutahedron P_n are indexed by up-rooted (or down-rooted) planar rooted leveled trees (PLTs). In particular, the vertices of P_n are indexed by the set $\wedge_n$ ($\vee_n$) of all up-rooted (down-rooted) planar binary trees with $n+1$ leaves and n levels. Since vertices of P_n are identified with permutations of $\underline{n}=\{1,2,\ldots,n\}$, the Bruhat partial ordering, generated by $a_1|\cdots|a_n<a_1|\cdots|a_{i+1}|a_i|\cdots|a_n$ if and only if $a_i<a_{i+1}$, induces natural poset structures on $\wedge_n$ and $\vee_n$.

In [22] we introduced the subposet $X^n_m\subseteq\wedge^{\times n}_m$, which indexes the vertices of the subcomplex $\Delta^{(n-1)}(P_m)\subseteq P^{\times n}_m$; an element $x\in X^n_m$ is expressed as a column matrix $x=[\hat{T}_1\cdots\hat{T}_n]^T$ of n up-rooted binary trees with $m+1$ leaves and m levels. Let $\curlywedge$ ($\curlyvee$) denote the up-rooted (down-rooted) 2-leaf corolla. Now if $a\in\wedge_n$, there exists a unique BTP Υ-factorization $a=a_1\cdots a_n$ such that a_j is a $1\times j$ row matrix over $\{\mathbf{1},\curlywedge\}$ containing the entry $\curlywedge$ exactly once. Thus by factoring each $\hat{T}_i$, we obtain the BTP Υ-factorization $x=x_1\cdots x_m$ in which x_j is an $n\times j$ matrix (see Example 6).

Let $\varrho(x,i)$ be the *lowest level* of $\hat{T}_i$ in which a branch is attached on the extreme left. Given positive integers $\alpha \leq \beta$, consider the set of *bitruncated elements*

$$X_m^n(\alpha,\beta) = \{{}_\alpha x_\beta = x_\alpha \cdots x_\beta \mid x = x_1 \cdots x_\alpha \cdots x_\beta \cdots x_m \in X_m^n,$$
$$\alpha = \min_{1\leq i\leq n} \varrho(x,i) \text{ and } \beta = \max_{1\leq i\leq n} \varrho(x,i)\}.$$

Define ${}_\alpha x_\beta < {}_{\alpha'} x'_{\beta'}$ if and only if $x < x'$ for some $x = x_1 \cdots x_\alpha \cdots x_\beta \cdots x_m$ and $x' = x'_1 \cdots x'_{\alpha'} \cdots x'_{\beta'} \cdots x'_m$ in X_m^n; then $X_m^n(\alpha,\beta)$ is the *poset of bitruncated elements*.

Now replace the entries $\mathbf{1}$ and $\curlywedge$ in each x_j with the integers 1 and 2, respectively. Then the $(m+1,m)$-row descent sequence $C_i = (\mathbf{x}_{i,1} = (2)\,, \mathbf{x}_{i,2}, \ldots, \mathbf{x}_{i,m})$ of $\hat{T}_i$ appears as the i^{th} rows of $x_1, \ldots, x_m$, and $\varrho = \varrho(x,i)$ is the largest integer such that $\mathbf{x}_{i,\varrho} = (2,1,...,1)$. Introduce the decorations $\mathring{\mathbf{x}}_{i,\varrho} = (\mathring{2}, \mathring{1}, ..., \mathring{1})$ and $\dot{\mathbf{x}}_{i,j} = (\dot{1}, 1, \ldots,\ 2, 1, \ldots)$ for $j > \varrho$, and obtain the corresponding $(m+1,m)$-*marked row descent sequence*

$$\hat{C}_i = (\mathbf{x}_{i,1}, \ldots, \mathbf{x}_{i,\varrho-1}, \mathring{\mathbf{x}}_{i,\varrho}, \dot{\mathbf{x}}_{i,\varrho+1}, \ldots, \dot{\mathbf{x}}_{i,m})\,.$$

The term $\mathring{\mathbf{x}}_{i,\varrho}$ represents the constant $1 \times \varrho$ matrix $\left[\mathfrak{f}_1^1 \cdots \mathfrak{f}_1^1\right]$. Let $\mathring{x}$ denote the matrix string $x = x_1 \cdots x_m$ with decorated integer entries; then the poset of *marked bitruncated elements* is

$$\mathring{X}_m^n(\alpha,\beta) = \{{}_\alpha \mathring{x}_\beta \mid_\alpha x_\beta \in X_m^n(\alpha,\beta)\}\,.$$

Note that $\varrho(x,i)$ is constant for all i if and only if $\alpha = \beta$, in which case $\mathring{X}_m^n(\alpha,\alpha)$ is a singleton set containing the $n \times \alpha$ matrix

$${}_\alpha \mathring{x}_\alpha = \begin{bmatrix} \mathring{\mathbf{x}}_{1,\alpha} \\ \vdots \\ \mathring{\mathbf{x}}_{n,\alpha} \end{bmatrix} = \begin{bmatrix} \mathring{2} & \mathring{1} & \cdots & \mathring{1} \\ \vdots & \vdots & & \vdots \\ \mathring{2} & \mathring{1} & \cdots & \mathring{1} \end{bmatrix},$$

which represents the constant matrix $\left[\mathfrak{f}_1^1\right]^{n\times\alpha}$.

Given ${}_\alpha \mathring{x}_\beta \in \mathring{X}_m^n(\alpha,\beta)$, consider the marked sequence of i^{th} rows

$$(\mathbf{x}_{i,\alpha}, \ldots, \mathbf{x}_{i,\varrho-1}, \mathring{\mathbf{x}}_{i,\varrho}, \dot{\mathbf{x}}_{i,\varrho+1}, \ldots, \dot{\mathbf{x}}_{i,\beta})\,.$$

Let $\mathbf{x}'_{i,k}$ denote the vector obtained from $\dot{\mathbf{x}}_{i,k}$ by deleting the marked entry $\dot{1}$, and form the (unmarked) row-descent sequence

$$(\mathbf{x}''_{i,\alpha}, ..., \mathbf{x}''_{i,\beta-1}) = (\mathbf{x}_{i,\alpha}, ..., \mathbf{x}_{i,\varrho-1}, \mathbf{x}'_{i,\varrho+1}, ..., \mathbf{x}'_{i,\beta}).$$

Then ${}_\alpha x''_{\beta-1}$ denotes the bitruncated element whose i^{th} rows are $\mathbf{x}''_{i,\alpha}, ..., \mathbf{x}''_{i,\beta-1}$.

Example 6. *Consider the following element $x \in X_6^2$ and its BTP factorization:*

$x =$



$$= \begin{bmatrix} \curlywedge \\ \curlywedge \end{bmatrix} \begin{bmatrix} \curlywedge\,\mathbf{1} \\ \curlywedge\,\mathbf{1} \end{bmatrix} \begin{bmatrix} \mathbf{1}\,\curlywedge\,\mathbf{1} \\ \mathbf{1}\,\curlywedge\,\mathbf{1} \end{bmatrix} \begin{bmatrix} \curlywedge\,\mathbf{1}\,\mathbf{1}\,\mathbf{1} \\ \mathbf{1}\,\mathbf{1}\,\curlywedge\,\mathbf{1} \end{bmatrix} \begin{bmatrix} \mathbf{1}\,\curlywedge\,\mathbf{1}\,\mathbf{1}\,\mathbf{1} \\ \mathbf{1}\,\curlywedge\,\mathbf{1}\,\mathbf{1}\,\mathbf{1} \end{bmatrix},$$

where the dotted lines indicate the lowest levels in which branches are attached on the extreme left ($\alpha = \varrho(x,2) = 2$ and $\beta = \varrho(x,1) = 4$). Then

$$\mathring{x} = \begin{bmatrix} 2 \\ 2 \end{bmatrix} \begin{bmatrix} 2 & 1 \\ \mathring{2} & \mathring{1} \end{bmatrix} \begin{bmatrix} 1 & 2 & 1 \\ \dot{1} & 2 & 1 \end{bmatrix} \begin{bmatrix} \mathring{2} & \mathring{1} & \mathring{1} & \mathring{1} \\ \dot{1} & 1 & 2 & 1 \end{bmatrix} \begin{bmatrix} \dot{1} & 2 & 1 & 1 & 1 \\ \dot{1} & 2 & 1 & 1 & 1 \end{bmatrix} \in \mathring{X}_6^2$$

and

$$ {}_2\mathring{x}_4 = \begin{bmatrix} 2 & 1 \\ \mathring{2} & \mathring{1} \end{bmatrix} \begin{bmatrix} 1 & 2 & 1 \\ \dot{1} & 2 & 1 \end{bmatrix} \begin{bmatrix} \mathring{2} & \mathring{1} & \mathring{1} & \mathring{1} \\ \dot{1} & 1 & 2 & 1 \end{bmatrix} \in \mathring{X}_6^2\,(2,4)\,. $$

The projections

$$ \left((2,1)\,,(1,2,1)\,,\left(\mathring{2},\mathring{1},\mathring{1},\mathring{1}\right)\right) \mapsto ((2,1)\,,(1,2,1)) $$

and

$$ \left(\left(\mathring{2},\mathring{1}\right),\left(\dot{1},2,1\right),\left(\dot{1},1,2,1\right)\right) \mapsto ((2,1)\,,(1,2,1)) $$

send ${}_2\mathring{x}_4$ *to*

$$ {}_2x_3'' = \begin{bmatrix} 2 & 1 \\ 2 & 1 \end{bmatrix} \begin{bmatrix} 1 & 2 & 1 \\ 1 & 2 & 1 \end{bmatrix}. $$

Dually, there is the subposet $Y_n^m \subseteq \vee_n^{\times m}$, which indexes the vertices of the subcomplex $\Delta^{(m-1)}(P_n) \subseteq P_n^{\times m}$; an element $y \in Y_n^m$ is expressed as a row matrix $y = [\check{T}_1 \cdots \check{T}_m]$ of m down-rooted binary trees with $n+1$ leaves and n levels. Now if $b \in \vee_n$, there exists a unique BTP $\boldsymbol{\Upsilon}$-factorization $b = b_n \cdots b_1$ such that b_i is a $i \times 1$ column matrix over $\{\mathbf{1}, \curlyvee\}$ containing the entry $\curlyvee$ exactly once. Thus by factoring each $\check{T}_j$, we obtain the (unique) BTP $\boldsymbol{\Upsilon}$-factorization $y = y_n \cdots y_1$ in which y_i is an $i \times m$ matrix (see Example 7).

Let $\varkappa(y,j)$ be the *highest level* of $\check{T}_j$ in which a branch is attached on the extreme right. Given positive integers $\epsilon \geq \delta$, consider the set of *bitruncated elements*

$$ \begin{gathered} Y_n^m(\epsilon,\delta) = \{{}_\epsilon y_\delta = y_\epsilon \cdots y_\delta \mid y = y_n \cdots y_\epsilon \cdots y_\delta \cdots y_1 \in Y_n^m, \\ \epsilon = \max_{1 \leq j \leq m} \varkappa(y,j) \text{ and } \delta = \min_{1 \leq j \leq m} \varkappa(y,j)\}. \end{gathered} $$

Define ${}_\epsilon y_\delta < {}_{\epsilon'} y'_{\delta'}$ if and only if $y < y'$ for some $y = y_n \cdots y_\epsilon \cdots y_\delta \cdots y_1$ and $y' = y'_n \cdots y'_{\epsilon'} \cdots y'_{\delta'} \cdots y'_1$ in Y_n^m; then $Y_n^m(\epsilon,\delta)$ is the *poset of bitruncated elements.*

Now replace the symbols $\mathbf{1}$ and $\curlyvee$ in each y_i with the integers 1 and 2, respectively. Then the $(n+1,n)$-column descent sequence $D_j = (\mathbf{y}_{n,j}, \ldots, \mathbf{y}_{2,j}, \mathbf{y}_{1,j} = (2))$ of $\check{T}_j$ appears as the j^{th} columns of $y_n, \ldots, y_1$, and $\varkappa = \varkappa(y,j)$ is the largest integer such that $\mathbf{y}_{\varkappa,j} = (1, ..., 1, 2)$. Introduce the decorations $\mathring{\mathbf{y}}_{\varkappa,j} = (\mathring{1}, ..., \mathring{1}, \mathring{2})^T$ and $\dot{\mathbf{y}}_{i,j} = \left(\ldots, 1, 2, \ldots, 1, \dot{1}\right)^T$ for $i > \varkappa$, and obtain the corresponding $(n+1,n)$-*marked column descent sequence*

$$ \check{D}_j = (\dot{\mathbf{y}}_{n,j}, \ldots, \dot{\mathbf{y}}_{\varkappa+1,j}, \mathring{\mathbf{y}}_{\varkappa,j}, \mathbf{y}_{\varkappa-1,j}, \ldots, \mathbf{y}_{1,j})\,. $$

The term $\mathring{\mathbf{y}}_{\varkappa,j}$ represents the constant $\varkappa \times 1$ matrix $\left[\mathfrak{f}_1^1 \cdots \mathfrak{f}_1^1\right]^T$. Let $\mathring{y}$ denote the matrix string $y = y_n \cdots y_1$ with decorated integer entries; then the poset of *marked bitruncated elements* is

$$ \mathring{Y}_n^m(\epsilon,\delta) = \{{}_\epsilon\mathring{y}_\delta \mid \ {}_\epsilon y_\delta \in Y_n^m(\epsilon,\delta)\}\,. $$

Note that $\varkappa(y,j)$ is constant for all j if and only if $\delta = \epsilon$, in which case $Y_n^m(\delta,\delta)$ is a singleton set containing the $\delta \times m$ matrix

$$ {}_\delta\mathring{y}_\delta = [\mathring{\mathbf{y}}_{\delta,1} \cdots \mathring{\mathbf{y}}_{\delta,m}] = \begin{bmatrix} \mathring{1} & \cdots & \mathring{1} \\ \vdots & & \vdots \\ \mathring{1} & \cdots & \mathring{1} \\ \mathring{2} & \cdots & \mathring{2} \end{bmatrix}, $$

which represents the constant matrix $\left[\mathfrak{f}_1^1\right]^{\delta \times m}$.

Given ${}_{\epsilon}\mathring{y}_{\delta} \in \mathring{Y}_n^m(\epsilon,\delta)$, consider the marked sequence of j^{th} columns

$$(\dot{\mathbf{y}}_{\epsilon,j},\ldots,\dot{\mathbf{y}}_{\varkappa+1,j},\mathring{\mathbf{y}}_{\varkappa,j},\mathbf{y}_{\varkappa-1,j},\ldots,\mathbf{y}_{\varkappa,j}).$$

Let $\mathbf{y}'_{k,j}$ denote the vector obtained from $\dot{\mathbf{y}}_{k,j}$ by deleting the marked entry $\dot{1}$, and form the (unmarked) column-descent sequence

$$\left(\mathbf{y}''_{\epsilon-1,j},...,\mathbf{y}''_{\delta,j}\right)=\left(\mathbf{y}'_{\epsilon-1,j},\ldots,\mathbf{y}'_{\varkappa+1,j},\mathbf{y}_{\varkappa-1,j},\ldots,\mathbf{y}_{\varkappa,j}\right).$$

Then ${}_{\epsilon-1}y''_{\delta}$ denotes the bitruncated element whose j^{th} columns are $\mathbf{y}''_{\epsilon-1,j},...,\mathbf{y}''_{\delta,j}$.

Example 7. *Consider the following element $y \in Y_6^2$ and its BTP factorization:*

$$y=$$



$$=\begin{bmatrix}\mathbf{1}\,\mathbf{1}\\ \Upsilon\Upsilon\\ \mathbf{1}\,\mathbf{1}\\ \mathbf{1}\,\mathbf{1}\\ \mathbf{1}\,\mathbf{1}\end{bmatrix}\begin{bmatrix}\mathbf{1}\,\mathbf{1}\\ \Upsilon\,\mathbf{1}\\ \mathbf{1}\,\mathbf{1}\\ \mathbf{1}\,\Upsilon\end{bmatrix}\begin{bmatrix}\mathbf{1}\,\mathbf{1}\\ \Upsilon\Upsilon\\ \mathbf{1}\,\mathbf{1}\end{bmatrix}\begin{bmatrix}\mathbf{1}\,\mathbf{1}\\ \Upsilon\Upsilon\end{bmatrix}[\Upsilon\Upsilon],$$

where the dotted lines indicate the highest levels in which branches are attached on the extreme right ($\epsilon=\varkappa(y,1)=4$ and $\delta=\varkappa(y,2)=2$). Then

$$\mathring{y}=\begin{bmatrix}1&1\\2&2\\1&1\\1&1\\\dot{1}&\dot{1}\end{bmatrix}\begin{bmatrix}1&\mathring{1}\\2&\mathring{1}\\1&\mathring{1}\\\dot{1}&\mathring{2}\end{bmatrix}\begin{bmatrix}1&1\\2&2\\\dot{1}&1\end{bmatrix}\begin{bmatrix}\mathring{1}&1\\\mathring{2}&2\end{bmatrix}\begin{bmatrix}2&2\end{bmatrix}\in\mathring{Y}_6^2$$

and

$${}_4\mathring{y}_2=\begin{bmatrix}1&\mathring{1}\\2&\mathring{1}\\1&\mathring{1}\\\dot{1}&\mathring{2}\end{bmatrix}\begin{bmatrix}1&1\\2&2\\\dot{1}&1\end{bmatrix}\begin{bmatrix}\mathring{1}&1\\\mathring{2}&2\end{bmatrix}\in\mathring{Y}_6^2(4,2).$$

The projections

$$\left(\begin{pmatrix}1\\2\\1\\\dot{1}\end{pmatrix},\begin{pmatrix}1\\2\\\dot{1}\end{pmatrix},\begin{pmatrix}\mathring{1}\\\mathring{2}\end{pmatrix}\right)\mapsto\left(\begin{pmatrix}1\\2\\1\end{pmatrix},\begin{pmatrix}1\\2\end{pmatrix}\right)$$

and

$$\left(\begin{pmatrix}\mathring{1}\\\mathring{1}\\\mathring{1}\\\mathring{2}\end{pmatrix},\begin{pmatrix}1\\2\\1\end{pmatrix},\begin{pmatrix}1\\2\end{pmatrix}\right)\mapsto\left(\begin{pmatrix}1\\2\\1\end{pmatrix},\begin{pmatrix}1\\2\end{pmatrix}\right)$$

send ${}_4\mathring{y}_2$ to

$${}_3y''_2=\begin{bmatrix}1&1\\2&2\\1&1\end{bmatrix}\begin{bmatrix}1&1\\2&2\end{bmatrix}.$$

Now extend the poset structures on these sets of bitruncated elements to

$$\mathring{X}_m^j(\alpha,\beta)\cup\mathring{Y}_m^i(\epsilon,\delta)$$

in the following way: Given ${}_{\alpha}\mathring{x}_{\beta} \in \mathring{X}^{j}_{m}(\alpha,\beta)$ and ${}_{\epsilon}\mathring{y}_{\delta} \in \mathring{Y}^{i}_{m}(\epsilon,\delta)$, define ${}_{\alpha}\mathring{x}_{\beta} \leq {}_{\epsilon}\mathring{y}_{\delta}$ if $\alpha \geq i$ and $j \leq \delta$, and ${}_{\epsilon}\mathring{y}_{\delta} \leq {}_{\alpha}\mathring{x}_{\beta}$ if $\alpha \leq i$ and $j \geq \delta$. Then $\alpha = i$ and $j = \beta$ implies $\alpha = \beta$ and $\epsilon = \delta$, in which case ${}_{\alpha}\mathring{x}_{\alpha} = {}_{\delta}\mathring{y}_{\delta}$. This equality reflects the correspondence

$$\begin{bmatrix} \mathring{2} & \mathring{1} & \cdots & \mathring{1} \\ \vdots & \vdots & & \vdots \\ \mathring{2} & \mathring{1} & \cdots & \mathring{1} \end{bmatrix}^{\delta\times\alpha} \leftrightarrow \begin{bmatrix} \mathfrak{f}^1_1 & \cdots & \mathfrak{f}^1_1 \\ \vdots & & \vdots \\ \mathfrak{f}^1_1 & \cdots & \mathfrak{f}^1_1 \end{bmatrix}^{\delta\times\alpha} \leftrightarrow \begin{bmatrix} \mathring{1} & \cdots & \mathring{1} \\ \vdots & & \vdots \\ \mathring{1} & \cdots & \mathring{1} \\ \mathring{2} & \cdots & \mathring{2} \end{bmatrix}^{\delta\times\alpha}.$$

Let $A = [a_{ij}]$ be an $(n+1)\times m$ matrix over $\{\mathbf{1},\curlywedge\}$, each row of which contains the entry $\curlywedge$ exactly once, and let $B = [b_{ij}]$ be an $n \times (m+1)$ matrix over $\{\mathbf{1},\curlyvee\}$, each column of which contains the entry $\curlyvee$ exactly once. Recall that a BTP (A,B) is an (i,j)-*edge pair* if $a_{ij} = a_{i+1,j} = \curlywedge$ and $b_{ij} = b_{i,j+1} = \curlyvee$. Note that (x_m, y_n) is the only potential edge pair in a matrix string $x_1 \cdots x_m y_n \cdots y_1 \in X^{n+1}_m \times Y^{m+1}_n$.

Given an (i,j)-edge pair (A,B), let A^{i*} and B^{*j} denote the matrices obtained by deleting the i^{th} row of A and the j^{th} column of B. If $c = C_1 \cdots C_k C_{k+1} \cdots C_r$ is a matrix string in which (C_k, C_{k+1}) is an (i,j)-edge pair, the (i,j)-*transposition of c in position k* is the matrix string

$$\mathcal{T}^k_{ij}(c) = C_1 \cdots C^{*j}_{k+1} C^{i*}_k \cdots C_r.$$

If (x_m, y_n) is an (i,j)-edge pair in $u = x_1 \cdots x_m y_n \cdots y_1 \in X^{n+1}_m \times Y^{m+1}_n$, then (x_{m-1}, y^{*j}_n) and (x^{i*}_m, y_{n-1}) are the potential edge pairs in $\mathcal{T}^m_{ij}(u)$. If (x_{m-1}, y^{*j}_n) is a (k,l)-edge pair, then (x_{m-2}, y^{*j*l}_n) is a potential edge pairs in

$$\mathcal{T}^{m-1}_{kl}\mathcal{T}^m_{ij}(u) = x_1 \cdots x_{m-2} y^{*j*l}_n x^{k*}_{m-1} x^{i*}_m y_{n-1} \cdots y_1,$$

and so on. Clearly, $\mathcal{T}^{k_t}_{i_t j_t} \cdots \mathcal{T}^{k_1}_{i_1 j_1}(u)$ uniquely determines a shuffle permutation $\sigma \in \Sigma_{m,n}$. On the other hand, (A,B) can be an (i,j)-edge pair for multiple values of i and j, in which case distinctly different compositions $\mathcal{T}^{k_t}_{i_t j_t} \cdots \mathcal{T}^{k_1}_{i_1 j_1}$ act on u and determine the same σ. Thus $\mathcal{T}_{\mathrm{Id}} := \mathrm{Id}$ and

$$\mathcal{T}_\sigma(u) := \left\{ \mathcal{T}^{k_t}_{i_t j_t} \cdots \mathcal{T}^{k_1}_{i_1 j_1}(u) \mid \mathcal{T}^{k_t}_{i_t j_t} \cdots \mathcal{T}^{k_1}_{i_1 j_1} \text{ determines } \sigma \in \Sigma_{m,n} \right\}.$$

Recall that $\mathcal{PP}_{n,m} = X^{n+1}_m \times Y^{m+1}_n \cup Z_{n,m}$, where

$$Z_{n,m} = \left\{ \mathcal{T}_\sigma(u) \mid u \in X^{n+1}_m \times Y^{m+1}_n \text{ and } \sigma \in \Sigma_{m,n} \smallsetminus \{\mathrm{Id}\} \right\}.$$

The poset $r\mathcal{PP}_{n,m}$ is built upon $\mathcal{PP}_{n,m}$. Given $u = x_1 \cdots x_m y_n \cdots y_1$ and $a = \mathcal{T}_\sigma(u)$, let $x^{\#}_i$ and $y^{\#}_j$ denote either x_i, y_j or their respective transpositions in a. Let

$$\mathcal{X}_{n,m} = \mathcal{X}'_{n,m} \cup \mathcal{X}''_{n,m}, \; m+n > 0, m, n \geq 0,$$

where

$$\mathcal{X}'_{n,m} = \left\{ (a, {}_{\alpha}\mathring{x}_{\alpha}) \in \bigcup_{\substack{1\leq\alpha\leq m+1 \\ 1\leq i\leq n+1}} \mathcal{PP}_{n,m} \times \mathring{X}^{i}_{m+1}(\alpha,\alpha) \;\middle|\; x^{\#}_{\alpha} \text{ has } i \text{ rows} \right\},$$

and

$$\mathcal{X}''_{n,m} = \left\{ (a, {}_{\alpha}\mathring{x}_{\beta}) \in \bigcup_{\substack{1\leq\alpha<\beta\leq m+1 \\ 1\leq i\leq n+1}} \mathcal{PP}_{n,m} \times \mathring{X}^{i}_{m+1}(\alpha,\beta) \;\middle|\; {}_{\alpha}x''_{\beta-1} \text{ is a substring of } a \right\}.$$

Dually, let

$$\mathcal{Y}_{n,m} = \mathcal{Y}'_{n,m} \cup \mathcal{Y}''_{n,m},\ \ m+n>0, m,n \geq 0,$$

where

$$\mathcal{Y}'_{n,m} = \left\{ (a,\ {}_{\delta}\mathring{y}_{\delta}) \in \bigcup_{\substack{1\leq\delta\leq n+1\\ 1\leq j\leq m+1}} \mathcal{PP}_{n,m} \times \mathring{Y}^{j}_{n+1}(\delta,\delta) \ \middle|\ y^{\#}_{\delta} \text{ has } j \text{ columns} \right\},$$

and

$$\mathcal{Y}''_{n,m} = \left\{ (a,\ {}_{\epsilon}\mathring{y}_{\delta}) \in \bigcup_{\substack{1\leq\delta<\epsilon\leq n+1\\ 1\leq j\leq m+1}} \mathcal{PP}_{n,m} \times \mathring{Y}^{j}_{n+1}(\epsilon,\delta) \ \middle|\ {}_{\epsilon-1}y''_{\delta} \text{ is a substring of } b \right\}.$$

Define

$$r\mathcal{PP}_{n,m} = \mathcal{X}_{n,m} \cup \mathcal{Y}_{n,m}$$

with the poset structure generated by $(a,b) \leq (a',b')$ for

- $a = a'$ and $b \leq b'$;
- $a \leq a'$ and $b = b'$ such that $u' = (\nu_x \times \nu_y)u$ for $\nu_x \in S^{\times n+1}_{\alpha-1} \times S^{\times n+1}_{\beta-\alpha} \times S^{\times n+1}_{m-\beta+1} \subset S^{\times n+1}_{m}$, $\nu_y \in S^{\times m+1}_{i-1} \times S^{\times m+1}_{n-i+1} \subset S^{\times m+1}_{n}$, $b \in \mathring{X}^{i}_{m+1}(\alpha,\beta), 1 \leq \alpha \leq \beta \leq m+1$, or $\nu_x \in S^{\times n+1}_{j-1} \times S^{\times n+1}_{m-j+1} \subset S^{\times n+1}_{m}$, $\nu_y \in S^{\times m+1}_{\delta-1} \times S^{\times m+1}_{\epsilon-\delta} \times S^{\times m+1}_{n-\epsilon+1} \subset S^{\times m+1}_{n}$, $b \in \mathring{Y}^{j}_{n+1}(\epsilon,\delta), 1 \leq \delta \leq \epsilon \leq n+1$ with convention that $S_0 \times S_k = S_k \times S_0 = S_k$, $k \geq 1$.

Note that when $\alpha = m+1$ in $\mathcal{X}'_{n,m}$ or $\delta = n+1$ in $\mathcal{Y}'_{n,m}$, in both cases $a \in X^{n+1}_{m} \times Y^{m+1}_{n} \subset \mathcal{P}P_{n,m}$ for $m,n \geq 1$; hence, the poset structure automatically identifies the subset

$$\mathcal{X}^{0}_{n,m} := \left(X^{n+1}_{m} \times Y^{m+1}_{n}\right) \times \mathring{X}^{n+1}_{m+1}(m+1,m+1) \subset \mathcal{X}_{n,m}$$

with the subset

$$\mathcal{Y}^{0}_{n,m} := \left(X^{n+1}_{m} \times Y^{m+1}_{n}\right) \times \mathring{Y}^{m+1}_{n+1}(n+1,n+1) \subset \mathcal{Y}_{n,m}.$$

Note also that $\mathcal{X}'_{m,0} = \mathcal{X}''_{m,0} = \mathcal{X}''_{0,m} = \mathcal{Y}'_{0,n} = \mathcal{X}''_{0,n} = \mathcal{Y}''_{n,0} = \varnothing$; thus $r\mathcal{PP}_{0,m} = \mathcal{X}'_{0,m}$ and $r\mathcal{PP}_{n,0} = \mathcal{Y}'_{n,0}$.

The poset $\mathcal{JJ}_{n+1,m+1}$ is the image of the quotient map $r\mathcal{PP}_{n,m} \to r\mathcal{PP}_{n,m}/\sim$ given by restricting the map $\mathcal{PP}_{*,*} \to \mathcal{KK}_{*+1,*+1}$ to left-hand factors, and $JJ_{n+1,m+1}$ is the geometric realization $|\mathcal{JJ}_{n+1,m+1}|$.

The poset structure of $r\mathcal{PP}$ corresponds to the "cylindrical poset" $\mathcal{KK} \times I$ in the following way. Given $a \in \mathcal{KK}_{n,m}$ and $t \geq m+n+1$, consider a partition

$$(a,b_1) < \cdots < (a,b_t)$$

of the interval $a \times I$, where $(a,b_1) = a \times 0$ and $(a,b_t) = a \times 1$. If a indexes a vertex $a_1|\cdots|a_{n+m} \in P_{n+m}$, and b_{t_α} denotes the single element of $\mathring{X}^{j}_{m}(\alpha,\alpha)$, then (a,b_{t_α}) indexes the vertex

$$a_1|\cdots|a_{n+\alpha-j}|m+n+1|a_{n+1+\alpha-j}|\cdots|a_{m+n} \in P_{m+n+1};$$

and in particular,

$$(a,b_1) \leftrightarrow a_1|\cdots|a_{m+n}|m+n+1 \text{ and } (a,b_t) \leftrightarrow m+n+1|a_1|\cdots|a_{m+n}.$$

Note that this correspondence agrees with the combinatorial representation of P_{m+n+1} as a subdivision of the cylinder $P_{m+n} \times P_2 = P_{m+n} \times I$ (see [20]), or equivalently, with the combinatorial join $P_{m+n+1} = P_{m+n} *_c P_1$ (see [22]). If in addition, $b_s \in \mathring{X}^j_{m+1}(\alpha,\beta)$ with $\beta = \alpha+1$, then (a, b_s) subdivides the interval $[(a, b_{t_{\alpha+1}}), (a, b_{t_\alpha})]$ (recall that $b_{t_{\alpha+1}} < b_{t_\alpha}$). In the octagon $JJ_{2,2}$ in Figure 5 and Example 8 below, we have

$$(1|2, b_1) < (1|2, b_2) < (1|2, b_3) < (1|2, b_4) < (1|2, b_5)$$

$$= 1|2|3 < v_1 < 1|3|2 < v_2 < 3|1|2,$$

where $v_1 \leftrightarrow b_2 \in \mathring{Y}^2_2(1,2)$ and $v_2 \leftrightarrow b_4 \in \mathring{X}^2_2(1,2)$; on the other hand,

$$(2|1, b_1) < (2|1, b_2) < (2|1, b_3)$$

$$= 2|1|3 < 2|3|1 < 3|2|1.$$

Now consider an element $(a,b) \in \mathcal{PP}_{n,m} \times \mathring{X}^j_{m+1}(\alpha,\beta) \subset r\mathcal{PP}_{n,m}$ and recall that $a = a_1 \cdots a_{n+m}$ is represented by a piecewise linear path in $\mathbb{N}^2$ with $m+n$ directed components. The element (a,b) is represented by a piecewise linear path in $\mathbb{N}^3$ of the form BCA with $m+n+1+\alpha-\beta$ directed components from $(m+1,1,0) \in \mathbb{N}^2 \times 0$ to $(1, n+1, 1) \in \mathbb{N}^2 \times 1$. The component A is represented by the path from $(m+1,1,0)$ to $(\beta, j, 0)$ in $\mathbb{N}^2 \times 0$ corresponding to $a_{n+\beta-j} \cdots a_{n+m}$; the component C is represented by the arrow $(\beta, j, 0) \to (\alpha, j, 1)$; and the component B is represented by the path from $(\alpha, j, 1)$ to $(1, n+1, 1)$ in $\mathbb{N}^2 \times 1$ corresponding to $a_1 \cdots a_{n+\alpha-j}$ (see Figure 1). Note that the arrow representing C is perpendicular to both integer lattices if and only if $\alpha = \beta$, in which case the path has $m+n+1$ directed components. The case of $(a,b) \in \mathcal{Y}_{n,m}$ is analogous with the arrow representing C lying in a vertical plane. For example, for $m = n = 1$, the singleton $(a,b) \in \mathcal{X}^0_{1,1} (= \mathcal{Y}^0_{1,1})$ is expressed by $DC'B$ in Figure 1 below.



Figure 1. $DEC < DC'B < C'''AB < C'''B'A'$.

Now suppose $w \in \mathcal{JJ}_{n+1,m+1}$ is the projection of $\tilde{w} = (a,b) \in r\mathcal{PP}_{n,m}$. Transform w into a 0-dimensional element of the A_∞-bialgebra morphism matrad $\mathcal{JJ}_\infty$ in the following way (see Example 8 below): If $\tilde{w} \in \mathcal{X}^0_{n,m}$ and $a = u \times v$, replace a by $u \cdot b \cdot v$; if $\tilde{w} \in \mathcal{X}'_{n,m}$, replace $a = \cdots x^{\#}_\alpha \cdots$ by $\cdots {}_\alpha\mathring{x}_\alpha \cdot x^{\#}_\alpha \cdots$; if $\tilde{w} \in \mathcal{X}''_{n,m}$ and $a = \cdots {}_\alpha\, \mathring{x}''_{\beta-1} \cdots$, replace ${}_\alpha x''_{\beta-1}$ by ${}_\alpha\mathring{x}_\beta = x_\alpha \cdots x_\beta$, if $\tilde{w} \in \mathcal{Y}'_{n,m}$, replace $a = \cdots y^{\#}_\delta \cdots$ by $\cdots y^{\#}_\delta \cdot {}_\delta\mathring{y}_\delta \cdots$; if $\tilde{w} \in \mathcal{Y}''_{n,m}$ and $a = \cdots {}_{\epsilon-1}\mathring{y}''_\delta \cdots$, replace ${}_{\epsilon-1}y''_\delta$ by ${}_\epsilon\mathring{y}_\delta = y_\epsilon \cdots y_\delta$. Now replace each row $(\mathring{2}, \mathring{1}, ..., \mathring{1})$ by $(\mathfrak{f}^1_1, \mathfrak{f}^1_1, ..., \mathfrak{f}^1_1)$ and each column

$(\mathring{1},...,\mathring{1},\mathring{2})^T$ by $(\mathfrak{f}_1^1,...,\mathfrak{f}_1^1,\mathfrak{f}_1^1)^T$; delete $\dot{1}$'s; replace each 1 by $\mathbf{1}$, replace each 2 in $x_i^\#$ by θ_2^1 and replace each 2 in $y_j^\#$ by θ_1^2. This transformation induces the bijection in (3.5) as in the absolute case.

Example 8. *The labels* v_1 *and* v_2 *are the midpoints of the edges* $1|23$ *and* $13|2$ *of* P_3*, respectively (see Figure 5).*

$$\begin{array}{rcll}
1|2|3 & \leftrightarrow & DEC & = \begin{bmatrix} \theta_2^1 \\ \theta_2^1 \end{bmatrix} \begin{bmatrix} \theta_1^2 & \theta_1^2 \end{bmatrix} \begin{bmatrix} \mathfrak{f}_1^1 & \mathfrak{f}_1^1 \end{bmatrix} \\
1|3|2 & \leftrightarrow & DC'B & = \begin{bmatrix} \theta_2^1 \\ \theta_2^1 \end{bmatrix} \begin{bmatrix} \mathfrak{f}_1^1 & \mathfrak{f}_1^1 \\ \mathfrak{f}_1^1 & \mathfrak{f}_1^1 \end{bmatrix} \begin{bmatrix} \theta_1^2 & \theta_1^2 \end{bmatrix} \\
3|1|2 & \leftrightarrow & C'''AB & = \begin{bmatrix} \mathfrak{f}_1^1 \\ \mathfrak{f}_1^1 \end{bmatrix} \begin{bmatrix} \theta_2^1 \\ \theta_2^1 \end{bmatrix} \begin{bmatrix} \theta_1^2 & \theta_1^2 \end{bmatrix} \\
2|1|3 & \leftrightarrow & E'D'C & = \left[\theta_1^2\right] \left[\theta_2^1\right] \begin{bmatrix} \mathfrak{f}_1^1 & \mathfrak{f}_1^1 \end{bmatrix} \\
2|3|1 & \leftrightarrow & E'C''A' & = \left[\theta_1^2\right] \left[\mathfrak{f}_1^1\right] \left[\theta_2^1\right] \\
3|2|1 & \leftrightarrow & C'''B'A' & = \begin{bmatrix} \mathfrak{f}_1^1 \\ \mathfrak{f}_1^1 \end{bmatrix} \left[\theta_1^2\right] \left[\theta_2^1\right] \\
v_1 & \leftrightarrow & \multicolumn{2}{l}{\begin{bmatrix} \theta_2^1 \\ \theta_2^1 \end{bmatrix} \begin{bmatrix} \left[\theta_1^2\right] \left[\mathfrak{f}_1^1\right] & \begin{bmatrix} \mathfrak{f}_1^1 \\ \mathfrak{f}_1^1 \end{bmatrix} \left[\theta_1^2\right] \end{bmatrix}} \\
v_2 & \leftrightarrow & \multicolumn{2}{l}{\begin{bmatrix} \left[\theta_2^1\right] \begin{bmatrix} \mathfrak{f}_1^1 & \mathfrak{f}_1^1 \end{bmatrix} \\ \left[\mathfrak{f}_1^1\right] \left[\theta_2^1\right] \end{bmatrix} \begin{bmatrix} \theta_1^2 & \theta_1^2 \end{bmatrix}.}
\end{array}$$

The bijection in (3.5) can be described on the codimension 1 level in the following way: The components $\left[\theta_{m+1}^{n+1}\right]\left[\mathfrak{f}_1^1\cdots\mathfrak{f}_1^1\right]$ and $\left[\mathfrak{f}_1^1\cdots\mathfrak{f}_1^1\right]\left[\theta_{m+1}^{n+1}\right]$ of $(\mathcal{J}\mathcal{J}_\infty)_{n+1,m+1}$ are assigned to the cells $KK_{n+1,m+1}\times 0$ and $KK_{n+1,m+1}\times 1$, which are the respective projections of $\underline{n+m} \mid n+m+1$ and $n+m+1 \mid \underline{n+m}$ in P_{m+n+1}, and labeled by the leaf sequences

$$\begin{matrix} \overbrace{1\cdots 1}^{n+1} \\ \underbrace{\mathring{1}\cdots\mathring{1}}_{m+1} \end{matrix} \quad \text{and} \quad \begin{matrix} \overbrace{\mathring{1}\cdots\mathring{1}}^{n+1} \\ \underbrace{1\cdots 1}_{m+1} \end{matrix}$$

respectively. The other components $A_p^{\mathbf{y}}R_{\mathbf{x}}^q$ and $Q_p^{\mathbf{y}}B_{\mathbf{x}}^q$ of $(\mathcal{J}\mathcal{J}_\infty)_{n+1,m+1}$ are assigned to the cells e_1 and e_2 of $JJ_{n+1,m+1}$ obtained by subdividing $K_{n+1,m+1}\times I$ in the following ways: Let $e_{(\mathbf{y},\mathbf{x})} = C|D$ be the codimension 1 cell of P_{m+n} defined in line (6.1) of [22]. Then

$$C|\left(D\cup\{m+n+1\}\right)\cup\left(C\cup\{m+n+1\}\right)|D = C|D\times I \subset P_{m+n}\times I \approx P_{m+n+1},$$

$$e_1 = (\vartheta_{n,m}\times 1)(C|D\cup\{m+n+1\}), \text{ and } e_2 = (\vartheta_{n,m}\times 1)\left(C\cup\{m+n+1\}|D\right).$$

The leaf sequences

$$\begin{matrix} \mathbf{y} \\ \mathring{\mathbf{x}} \end{matrix} \quad \text{and} \quad \begin{matrix} \mathring{\mathbf{y}} \\ \mathbf{x} \end{matrix}$$

label e_1 and e_2, respectively (see Figures 2-6 below).

Unlike the 3-dimensional biassociahedra, certain 2-cells of the 3-dimensional bimultiplihedra are defined in terms of Δ_P. When constructing $JJ_{2,3}$, for example, we use $\Delta_P(P_3)$ to subdivide the cell 124|3 and label the faces in a manner similar to the labeling on $KK_{2,3}$ (see [22], Figure 19). Furthermore, it is convenient to think of $JJ_{2,3}$ as a subdivision of the cylinder $KK_{2,3} \times I$; then 2-faces of $(\mathcal{JJ}_\infty)_{2,3}$ are represented by a path $(3,1,0) \to (s,t,\epsilon) \to (1,2,1)$ in $\mathbb{N}^3$ with $\epsilon = 0, 1$. For example, the paths $(3,1,0) \to (2,2,1) \to (1,2,1)$ and $(3,1,0) \to (2,2,0) \to (1,2,1)$ represent the 2-faces $e_1 = 1|234$ and $e_2 = 14|23$, respectively. Finally, we remark that a general codimension 1 face of $(\mathcal{JJ}_\infty)_{n,m}$ is detected by the pair of leaf sequences $(\mathbf{x}, \mathbf{y})$ and the corresponding path.

While $JJ_{1,n}$ and $JJ_{n,1}$ are combinatorially isomorphic to the multiplihedron J_n, their distinct combinatorial representations are related by the cellular isomorphism $JJ_{1,n} \to JJ_{n,1}$ induced by the reversing map $\chi : a_1|\cdots|a_n \mapsto a_n|\cdots|a_1$ on P_n. Let $\pi_1 : P_n \to J_n$ and $\pi_2 : P_n \to J_n$ be the respective cellular projections $P_n = |r\mathcal{PP}_{0,n-1}| \to JJ_{1,n}$ and $P_n = |r\mathcal{PP}_{n-1,0}| \to JJ_{n,1}$, $n \geq 2$; then $\pi_1 = \pi_2 \circ \chi$.

Example 9. *The bijection (3.5) is represented on the vertices of* J_4 *in* $\pi_1(2|134 \cup 24|13)$ *as follows:*

$$\begin{array}{ccc}
[\theta_2^1]\ [\mathbf{1}\ \theta_2^1]\ [\theta_2^1\ \mathbf{1}\ \mathbf{1}]\ [\mathfrak{f}_1^1\ \mathfrak{f}_1^1\ \mathfrak{f}_1^1\ \mathfrak{f}_1^1] & \sim & [\theta_2^1]\ [\theta_2^1\ \mathbf{1}]\ [\mathbf{1}\ \mathbf{1}\ \theta_2^1]\ [\mathfrak{f}_1^1\ \mathfrak{f}_1^1\ \mathfrak{f}_1^1\ \mathfrak{f}_1^1] \\
\updownarrow & & \updownarrow \\
((2),(12),(211),(\mathring{2}\mathring{1}\mathring{1}\mathring{1})) & \sim & ((2),(21),(112),(\mathring{2}\mathring{1}\mathring{1}\mathring{1})) \\
\updownarrow & & \updownarrow \\
2|1|3|4 & \overset{\pi_1}{\sim} & 2|3|1|4
\end{array}$$

$$\begin{array}{ccc}
[\theta_2^1]\ [\mathbf{1}\ \theta_2^1]\ [\mathfrak{f}_1^1\ \mathfrak{f}_1^1\ \mathfrak{f}_1^1]\ [\theta_2^1\ \mathbf{1}\ \mathbf{1}] & & [\theta_2^1]\ [\theta_2^1\ \mathbf{1}]\ [\mathfrak{f}_1^1\ \mathfrak{f}_1^1\ \mathfrak{f}_1^1]\ [\mathbf{1}\ \mathbf{1}\theta_2^1] \\
\updownarrow & & \updownarrow \\
((2),(12),(\mathring{2}\mathring{1}\mathring{1}),(\dot{1}211)) & & ((2),(21),(\mathring{2}\mathring{1}\mathring{1}),(\dot{1}112)) \\
\updownarrow & & \updownarrow \\
2|1|4|3 & & 2|3|4|1
\end{array}$$

$$\begin{array}{ccc}
[\theta_2^1]\ [\mathfrak{f}_1^1\ \mathfrak{f}_1^1]\ [\mathbf{1}\ \theta_2^1]\ [\theta_2^1\ \mathbf{1}\ \mathbf{1}] & \sim & [\theta_2^1]\ [\mathfrak{f}_1^1\ \mathfrak{f}_1^1]\ [\theta_2^1\ \mathbf{1}]\ [\mathbf{1}\ \mathbf{1}\ \theta_2^1] \\
\updownarrow & & \updownarrow \\
((2),(\mathring{2}\mathring{1}),(\dot{1}12),(\dot{1}211)) & \sim & ((2),(\mathring{2}\mathring{1}),(\dot{1}21),(\dot{1}112)) \\
\updownarrow & & \updownarrow \\
2|4|1|3 & \overset{\pi_1}{\sim} & 2|4|3|1
\end{array}$$

$$\begin{array}{ccc}
[\mathfrak{f}_1^1]\ [\theta_2^1]\ [\mathbf{1}\ \theta_2^1]\ [\theta_2^1\ \mathbf{1}\ \mathbf{1}] & \sim & [\mathfrak{f}_1^1]\ [\theta_2^1]\ [\theta_2^1\ \mathbf{1}]\ [\mathbf{1}\ \mathbf{1}\ \theta_2^1] \\
\updownarrow & & \updownarrow \\
((\mathring{2}),(\dot{1}2),(\dot{1}12),(\dot{1}211)) & \sim & ((\mathring{2}),(\dot{1}2),(\dot{1}21),(\dot{1}112)) \\
\updownarrow & & \updownarrow \\
4|2|1|3 & \overset{\pi_1}{\sim} & 4|2|3|1.
\end{array}$$

Note that the 2-cell 24|13 *and the edge* 2|13|4 *of* P_4 *degenerate under* π_1*, whereas the 2-cell* 13|24 *and the edge* 4|13|2 *of* P_4 *degenerate under* π_2*. For comparison, the cellular map* $\pi : P_n \to J_n$ *defined in* [20] *differs from both* π_1 *and* π_2*: The 2-cell* 24|13 *and the edge* 1|24|3 *of* P_4 *degenerate under* π*.*


1
$\begin{smallmatrix}\mathring{1}\\ \mathring{1}\end{smallmatrix}$

Figure 2. The point $JJ_{1,1}$.


1|2
2|1
$\begin{smallmatrix}2\\ \mathring{1}\end{smallmatrix}$
$\begin{smallmatrix}\mathring{1}\mathring{1}\\ 1\end{smallmatrix}$

Figure 3. The interval $JJ_{2,1}$.


1|2
2|1
$\begin{smallmatrix}1\\ \mathring{1}\mathring{1}\end{smallmatrix}$
$\begin{smallmatrix}\mathring{1}\\ 2\end{smallmatrix}$

Figure 4. The interval $JJ_{1,2}$.


3|12
13|2
$\begin{smallmatrix}\mathring{1}\mathring{1}\\ 11\end{smallmatrix}$
$\begin{smallmatrix}\mathring{1}\mathring{1}\\ 2\end{smallmatrix}$
$\begin{smallmatrix}\mathring{2}\\ 2\end{smallmatrix}$
23|1
1|23
$\begin{smallmatrix}11\\ \mathring{1}\mathring{1}\end{smallmatrix}$
$\begin{smallmatrix}2\\ \mathring{2}\end{smallmatrix}$
2|13
$\begin{smallmatrix}2\\ \mathring{1}\mathring{1}\end{smallmatrix}$
12|3

Figure 5. The octagon $JJ_{2,2}$ as a subdivision of P_3.

Combinatorial data For $JJ_{2,3}$:

123|4	↔	$\begin{smallmatrix}2\\ \mathring{1}\mathring{1}\mathring{1}\end{smallmatrix}$	=	$\theta_3^2/\mathfrak{f}_1^1\mathfrak{f}_1^1\mathfrak{f}_1^1$	1|234	↔	$\begin{smallmatrix}11\\ \mathring{2}\mathring{1}\end{smallmatrix}$	=	$\theta_2^1\theta_2^1/\mathfrak{f}_2^2\left(\mathfrak{f}_1^1\mathfrak{f}_1^1/\theta_1^2\right)$
4|123	↔	$\begin{smallmatrix}\mathring{1}\mathring{1}\\ 3\end{smallmatrix}$	=	$\mathfrak{f}_1^1\mathfrak{f}_1^1/\theta_3^2$	14|23	↔	$\begin{smallmatrix}\mathring{1}\mathring{1}\\ 21\end{smallmatrix}$	=	$[(\mathfrak{f}_1^1/\theta_2^1)\mathfrak{f}_2^1+\mathfrak{f}_2^1\left(\theta_2^1/\mathfrak{f}_1^1\mathfrak{f}_1^1\right)]/\theta_2^2\theta_1^2$
13|24	↔	$\begin{smallmatrix}2\\ \mathring{2}\mathring{1}\end{smallmatrix}$	=	$\theta_2^2/\mathfrak{f}_2^1\mathfrak{f}_1^1$	2|134	↔	$\begin{smallmatrix}11\\ \mathring{1}\mathring{2}\end{smallmatrix}$	=	$\theta_2^1\theta_2^1/(\theta_1^2/\mathfrak{f}_1^1)\mathfrak{f}_2^2$
134|2	↔	$\begin{smallmatrix}\mathring{2}\\ 21\end{smallmatrix}$	=	$\mathfrak{f}_2^2/\theta_2^1\theta_1^1$	24|13	↔	$\begin{smallmatrix}\mathring{1}\mathring{1}\\ 12\end{smallmatrix}$	=	$[(\mathfrak{f}_1^1/\theta_2^1)\mathfrak{f}_2^1+\mathfrak{f}_2^1\left(\theta_2^1/\mathfrak{f}_1^1\mathfrak{f}_1^1\right)]/\theta_1^2\theta_2^2$
3|124	↔	$\begin{smallmatrix}2\\ \mathring{3}\end{smallmatrix}$	=	$\theta_1^2/\mathfrak{f}_3^1$	23|14	↔	$\begin{smallmatrix}2\\ \mathring{1}\mathring{2}\end{smallmatrix}$	=	$\theta_2^2/\mathfrak{f}_1^1\mathfrak{f}_2^1$
34|12	↔	$\begin{smallmatrix}\mathring{2}\\ 3\end{smallmatrix}$	=	$\mathfrak{f}_1^2/\theta_3^1$	234|1	↔	$\begin{smallmatrix}\mathring{2}\\ 12\end{smallmatrix}$	=	$\mathfrak{f}_2^2/\theta_1^1\theta_2^1$

$$\begin{aligned}
12|34 \quad \leftrightarrow \quad \substack{11\\ \mathring{1}\mathring{1}\mathring{1}} \quad &= \quad [\left(\theta_2^1/\theta_2^1\theta_1^1\right)\theta_3^1+\theta_3^1\left(\theta_2^1/\theta_1^1\theta_2^1\right)]/\\
&\quad\ [(\theta_1^2/\mathfrak{f}_1^1)(\theta_1^2/\mathfrak{f}_1^1)\mathfrak{f}_1^2+\ (\theta_1^2/\mathfrak{f}_1^1)\mathfrak{f}_1^2(\mathfrak{f}_1^1\mathfrak{f}_1^1/\theta_1^2)+\mathfrak{f}_1^2(\mathfrak{f}_1^1\mathfrak{f}_1^1/\theta_1^2)(\mathfrak{f}_1^1\mathfrak{f}_1^1/\theta_1^2)]\\
124|3 \quad \leftrightarrow \quad \substack{\mathring{1}\mathring{1}\\ 111} \quad &= \quad [(\theta_2^1/\theta_2^1\theta_1^1/\mathfrak{f}_1^1\mathfrak{f}_1^1\mathfrak{f}_1^1)\mathfrak{f}_3^1+\mathfrak{f}_3^1(\mathfrak{f}_1^1/\theta_2^1/\theta_1^1\theta_2^1)+(\theta_3^1/\mathfrak{f}_1^1\mathfrak{f}_1^1\mathfrak{f}_1^1)(\theta_2^1/\mathfrak{f}_1^1\mathfrak{f}_2^1)\\
&\quad\ +(\theta_3^1/\mathfrak{f}_1^1\mathfrak{f}_1^1\mathfrak{f}_1^1)(\mathfrak{f}_2^1/\theta_1^1\theta_2^1)+(\theta_2^1/\mathfrak{f}_1^1\mathfrak{f}_2^1)(\mathfrak{f}_2^1/\theta_1^1\theta_2^1)-(\theta_2^1/\mathfrak{f}_2^1\mathfrak{f}_1^1)(\mathfrak{f}_2^1/\theta_2^1\theta_1^1)\\
&\quad\ -(\theta_2^1/\mathfrak{f}_2^1\mathfrak{f}_1^1)(\mathfrak{f}_1^1/\theta_3^1)-(\mathfrak{f}_2^1/\theta_2^1\theta_1^1)(\mathfrak{f}_1^1/\theta_3^1)]/\,\theta_1^2\theta_1^2\theta_1^2
\end{aligned}$$



Figure 6. The bimultiplihedron $JJ_{2,3}$ as a subdivision of P_4.

5. Morphisms Defined

In this section we define the morphisms of A_∞-bialgebras. Our approach is to think of a morphism of A_∞-(co)algebras as an $\mathcal{A}_\infty$-bimodule; and analogously, to think of a morphism of A_∞-bialgebras as an $\mathcal{H}_\infty$-bimodule. But before we begin, we mention three settings in which A_∞-bialgebras naturally appear:

(1) Let X be a space and let $S_*(X)$ denote the simplicial singular chain complex of X. Then the cobar construction $\Omega S_*(X)$ is a DG bialgebra with coassociative coproduct [1], [5], [12]. While it is highly improbably that the double cobar construction $\Omega^2 S_*(X)$ admits a coassociative coproduct, there is an A_∞-coalgebra structure on $\Omega^2 S_*(X)$ that is compatible with the product, and $\Omega^2 S_*(X)$ is an A_∞-bialgebra.

(2) Let H be a graded bialgebra with nontrivial product and coproduct, and let $\rho : RH \longrightarrow H$ be a (bigraded) multiplicative resolution. Since RH cannot be simultaneously free and cofree, it is difficult to introduce a strictly coassociative coproduct on RH in such a way that ρ is a bialgebra map. However, there is an A_∞-bialgebra structure on RH such that ρ is a morphism of A_∞-bialgebras.

(3) If A is an A_∞-bialgebra over a field and $g : H(A) \to A$ is a cycle selecting homomorphism on homology, there is an A_∞-bialgebra structure on $H(A)$ that

is unique up to isomorphism, and a morphism of A_∞-bialgebras $G : H(A) \Rightarrow A$ extending g (see Theorem 2).

Recall the following equivalent definitions of an A_∞-bialgebra:

Definition 18. *A graded R-module A together with an element*

$$\omega = \{\omega_m^n \in Hom^{m+n-3}(A^{\otimes m}, A^{\otimes n})\}_{m,n\geq 1} \in U_A$$

*is an A_∞**-bialgebra** if either*

(i) $\omega \circledcirc \omega = 0$ or
(ii) the map $\theta_m^n \mapsto \omega_m^n$ extends to a map $\mathcal{H}_\infty \to U_A$ of matrads.

There is an operation $\ominus : U_B \times U_{A,B} \times U_A \to U_{A,B}$ analogous to $\circledcirc$, which allows us to define a morphism of A_∞-bialgebras in two equivalent ways. Given DG R-modules (DGMs) (A, d_A) and (B, d_B), let $d_A : TA \to TA$ and $d_B : TB \to TB$ be the free linear extensions of d_A and d_B, and let ∇ be the induced Hom differential on $U_{A,B}$, i.e., for $f \in U_{A,B}$ define $\nabla f = d_B \circ f - (-1)^{|f|} f \circ d_A$. Given $(\zeta, f, \eta) \in U_B \times U_{A,B} \times U_A$ and $m, n \geq 1$, obtain $X^{n,m} + Y^{n,m} \in (U_{A,B})_{n,m}$ by replacing $(\theta, \mathfrak{f}, \theta)$ in the right-hand side of formula (3.4) with (ζ, f, η). Then define

$$\ominus(\zeta, f, \eta) = \{X^{n,m}\}_{m,n\geq 1} + \{Y^{n,m}\}_{m,n\geq 1} - \nabla f.$$

Definition 19. *Let (A, ω_A) and (B, ω_B) be A_∞-bialgebras. An element*

$$G = \{g_m^n \in Hom^{m+n-2}(A^{\otimes m}, B^{\otimes n})\}_{m,n\geq 1} \in U_{A,B}$$

*is an A_∞-**bialgebra morphism from** A **to** B if either*

(i) $\ominus(\omega_B, G, \omega_A) = 0$ or
(ii) the map $\mathfrak{f}_m^n \mapsto g_m^n$ extends to a map $\mathcal{J}\mathcal{J}_\infty \to U_{A,B}$ of relative matrads.

*The symbol $G : A \Rightarrow B$ denotes an A_∞-bialgebra morphism G from A to B. An A_∞-bialgebra morphism $\Phi = \{\phi_m^n\}_{m,n\geq 1} : A \Rightarrow B$ is an **isomorphism** if $\phi_1^1 : A \to B$ is an isomorphism of underlying modules.*

6. Transfer of A_∞-Structure

If A is a free DGM, B is an A_∞-algebra, and $g : A \to B$ is a homology isomorphism (weak equivalence) with a right-homotopy inverse, the Basic Perturbation Lemma (BPL) transfers the A_∞-algebra structure from B to A (see [9], [15], for example). When B is an A_∞-bialgebra, Theorem 1 generalizes the BPL in two directions:

1. We transfer the A_∞-bialgebra structure from B to A.
2. Our transfer algorithm requires neither freeness nor the existence of a right-homotopy inverse.

Note that generalization (2) formulates the classical transfer of A_∞-algebra structure in maximal generality (see Remark 2).

A chain map $g : A \to B$ of DGMs induces a cochain map $\tilde{g} : U_A \to U_{A,B}$ defined on $u \in Hom\,(A^{\otimes m}, A^{\otimes n})$ by $\tilde{g}\,(u) = g^{\otimes n} u$, which is a homology isomorphism if the hypotheses of our next proposition are satisfied (the proof is left to the reader):

Proposition 1. *Let (A, d_A) and (B, d_B) be DGMs, let $g : A \to B$ be a chain map that is also a homology isomorphism. Then $\tilde{g} : U_A \to U_{A,B}$ is a homology isomorphism whenever either of the following conditions holds:*

(i) A is free as an R-module; or

(ii) for each $n \geq 1$*, there is a DGM* $X(n)$ *and a splitting* $B^{\otimes n} = A^{\otimes n} \oplus X(n)$ *as a chain complex such that* $H^*Hom\left(A^{\otimes k}, X(n)\right) = 0$ *for all* $k \geq 1$*.*

Since the hypotheses of Proposition 1 are satisfied when R is a field, there is the following generalization of the BPL:

Theorem 1 (**The Transfer**). *Let* (A, d_A) *be a DGM, let* (B, d_B, ω_B) *be an* A_∞*-bialgebra, and let* $g : A \to B$ *be a chain map that is also a homology isomorphism. If* $\tilde{g} : \mathcal{E}nd_{TA} \to U_{A,B}$ *is a homology isomorphism, then*

(i) (Existence) g *induces an* A_∞*-bialgebra structure* $\omega_A = \{\omega_A^{n,m}\}$ *on* A *and extends to a map* $G = \{g_m^n \mid g_1^1 = g\} : A \Rightarrow B$ *of* A_∞*-bialgebras.*

(ii) (Uniqueness) (ω_A, G) *is unique up to isomorphism, i.e., if* (ω_A, G) *and* $\left(\bar{\omega}_A, \bar{G}\right)$ *are induced by chain homotopic maps* g *and* $\bar{g}$*, there is an isomorphism* $\Phi : (A, \bar{\omega}_A) \Rightarrow (A, \omega_A)$ *and a chain homotopy* $T : \bar{G} \simeq G \circ \Phi$*.*

Proof. We obtain the desired structures by simultaneously constructing a map of matrads $\alpha_A : C_*(KK) \to U_A$ and a map of relative matrads $\beta : C_*(JJ) \to U_{A,B}$. Thinking of $JJ_{n,m}$ as a subdivision of the cylinder $KK_{n,m} \times I$, identify the top dimensional cells of $KK_{n,m}$ and $JJ_{n,m}$ with θ_m^n and $\mathfrak{f}_m^n$, and the faces $KK_{n,m} \times 0$ and $KK_{n,m} \times 1$ of $JJ_{n,m}$ with $\theta_m^n \left(\mathfrak{f}_1^1\right)^{\otimes m}$ and $\left(\mathfrak{f}_1^1\right)^{\otimes n} \theta_m^n$, respectively. By hypothesis, there is a map of matrads $\alpha_B : C_*(KK) \to U_B$ such that $\alpha_B(\theta_m^n) = \omega_B^{n,m}$.

To initialize the induction, define $\beta : C_*(JJ_{1,1}) \to Hom^0(A, B)$ by $\beta\left(\mathfrak{f}_1^1\right) = g_1^1 = g$ and extend β to $C_*(JJ_{1,2}) \to Hom^1\left(A^{\otimes 2}, B\right)$ and $C_*(JJ_{2,1}) \to Hom^1\left(A, B^{\otimes 2}\right)$ in the following way: On the vertices $\theta_2^1\left(\mathfrak{f}_1^1 \otimes \mathfrak{f}_1^1\right) \in JJ_{1,2}$ and $\theta_1^2 \mathfrak{f}_1^1 \in JJ_{2,1}$ define $\beta\left(\theta_2^1\left(\mathfrak{f}_1^1 \otimes \mathfrak{f}_1^1\right)\right) = \omega_B^{1,2}(g \otimes g)$ and $\beta\left(\theta_1^2 \mathfrak{f}_1^1\right) = \omega_B^{2,1} g$. Since $\omega_B^{1,2}(g \otimes g)$ and $\omega_B^{2,1} g$ are ∇-cocycles and $\tilde{g}_*$ is an isomorphism, there exist cocycles $\omega_A^{1,2}$ and $\omega_A^{2,1}$ in U_A such that

$$\tilde{g}_*[\omega_A^{1,2}] = [\omega_B^{1,2}(g \otimes g)] \text{ and } \tilde{g}_*[\omega_A^{2,1}] = [\omega_B^{2,1} g].$$

Thus $\left[\omega_B^{1,2}(g \otimes g) - g\omega_A^{1,2}\right] = \left[\omega_B^{2,1} g - (g \otimes g)\,\omega_A^{2,1}\right] = 0$, and there exist cochains g_2^1 and g_1^2 in $U_{A,B}$ such that

$$\nabla g_2^1 = \omega_B^{1,2}(g \otimes g) - g\omega_A^{1,2} \text{ and } \nabla g_1^2 = \omega_B^{2,1} g - (g \otimes g)\,\omega_A^{2,1}.$$

For $m = 1, 2$ and $n = 3 - m$, define $\alpha_A : C_*(KK_{n,m}) \to Hom\left(A^{\otimes m}, A^{\otimes n}\right)$ by $\alpha_A(\theta_m^n) = \omega_A^{n,m}$ and define $\beta : C_*(JJ_{n,m}) \to Hom\left(A^{\otimes m}, B^{\otimes n}\right)$ by

$$\begin{array}{rcll} \beta(\mathfrak{f}_m^n) & = & g_m^n & \\ \beta(\mathfrak{f}_1^1\,\theta_2^1) & = & g\,\omega_A^{1,2} & (m=2) \\ \beta((\mathfrak{f}_1^1 \otimes \mathfrak{f}_1^1)\,\theta_1^2) & = & (g \otimes g)\,\omega_A^{2,1} & (m=1). \end{array}$$

Inductively, given (m, n), $m + n \geq 4$, assume that for $i + j < m + n$ there exists a map of matrads $\alpha_A : C_*(KK_{j,i}) \to Hom\left(A^{\otimes i}, A^{\otimes j}\right)$ and a map of relative matrads $\beta : C_*(JJ_{j,i}) \to Hom\left(A^{\otimes i}, B^{\otimes j}\right)$ such that $\alpha_A(\theta_i^j) = \omega_A^{j,i}$ and $\beta(\mathfrak{f}_i^j) = g_i^j$. Thus we are given chain maps $\alpha_A : C_*(\partial KK_{n,m}) \to Hom\left(A^{\otimes m}, A^{\otimes n}\right)$ and $\beta : C_*(\partial JJ_{n,m} \smallsetminus \operatorname{int} KK_{n,m} \times 1) \to Hom\left(A^{\otimes m}, B^{\otimes n}\right)$; we wish to extend α_A to the top cell θ_m^n of $KK_{n,m}$ and β to the codimension 1 cell $\left(\mathfrak{f}_1^1\right)^{\otimes n} \theta_m^n$ and the top cell $\mathfrak{f}_m^n$ of $JJ_{n,m}$. Since α_A is a map of matrads, the components of the cocycle

$$z = \alpha_A\left(C_*(\partial KK_{n,m})\right) \in Hom^{m+n-4}\left(A^{\otimes m}, A^{\otimes n}\right)$$

are expressed in terms of $\omega_A^{j,i}$ with $i+j<m+n$; similarly, since β is a map of relative matrads, the components of the cochain

$$\varphi=\beta\left(C_*(\partial JJ_{n,m}\smallsetminus\operatorname{int}KK_{n,m}\times 1)\right)\in Hom^{m+n-3}\left(A^{\otimes m},B^{\otimes n}\right)$$

are expressed in terms of ω_B, $\omega_A^{j,i}$ and g_i^j with $i+j<m+n$. Clearly $\tilde{g}\left(z\right)=\nabla\varphi$; and $[z]=[0]$ since $\tilde{g}$ is a homology isomorphism. Now choose a cochain $b\in Hom^{m+n-3}\left(A^{\otimes m},A^{\otimes n}\right)$ such that $\nabla b=z$. Then

$$\nabla\left(\tilde{g}\left(b\right)-\varphi\right)=\nabla\tilde{g}\left(b\right)-\tilde{g}\left(z\right)=0.$$

Choose a class representative $u\in\tilde{g}_*^{-1}\left[\tilde{g}\left(b\right)-\varphi\right]$, set $\omega_A^{n,m}=b-u$, and define $\alpha_A\left(\theta_m^n\right)=\omega_A^{n,m}$. Then $\left[\tilde{g}\left(\omega_A^{n,m}\right)-\varphi\right]=\left[\tilde{g}\left(b-u\right)-\varphi\right]=\left[\tilde{g}\left(b\right)-\varphi\right]-\left[\tilde{g}\left(u\right)\right]=\left[0\right].$ Choose a cochain $g_m^n\in Hom^{m+n-2}\left(A^{\otimes m},B^{\otimes n}\right)$ such that

$$\nabla g_m^n=g^{\otimes n}\omega_A^{n,m}-\varphi,$$

and define $\beta\left(\mathfrak{f}_m^n\right)=g_m^n$. To extend β as a map of relative matrads, define $\beta\left(\left(\mathfrak{f}_1^1\right)^{\otimes n}\theta_m^n\right)=g^{\otimes n}\omega_A^{n,m}$. Passing to the limit we obtain the desired maps α_A and β.

Furthermore, if chain maps $\bar{\alpha}_A$ and $\bar{\beta}$ are defined in terms of different choices, beginning with a chain map $\bar{g}$ chain homotopic to g, let $\bar{\omega}_A=\operatorname{Im}\bar{\alpha}_A$ and $\bar{G}=\operatorname{Im}\bar{\beta}$. There is an inductively defined isomorphism $\Phi=\sum\phi_m^n:(A,\bar{\omega}_A)\Rightarrow(A,\omega_A)$ with $\phi_1^1=\mathbf{1}$, and a chain homotopy $T:\tilde{G}\simeq G\circ\Phi$. To initialize the induction, set $\phi_1^1=\mathbf{1}$ and note that

$$\nabla g_2^1=g\omega_A^{1,2}-\omega_B^{1,2}(g\otimes g)\text{ and }\nabla\bar{g}_2^1=\bar{g}\bar{\omega}_A^{1,2}-\omega_B^{1,2}(\bar{g}\otimes\bar{g}).$$

Let $s:\bar{g}\simeq g$; then $c_2^1=\omega_B^{1,2}\left(s\otimes g+\bar{g}\otimes s\right)$ satisfies

$$\nabla c_2^1=\omega_B^{1,2}(g\otimes g)-\omega_B^{1,2}(\bar{g}\otimes\bar{g}).$$

Hence

$$\nabla(g_2^1-\bar{g}_2^1+c_2^1)=g\omega_A^{1,2}-\bar{g}\bar{\omega}_A^{1,2}$$

and

$$\bar{g}(\omega_A^{1,2}-\bar{\omega}_A^{1,2})=\nabla(g_2^1-\bar{g}_2^1+c_2^1-s\,\omega_A^{1,2}).$$

Consequently, there is $\phi_2^1:A^{\otimes 2}\to A$ such that $\nabla\phi_2^1=\omega_A^{1,2}-\bar{\omega}_A^{1,2}$; and, as above, ϕ_2^1 may be chosen so that $\bar{g}\phi_2^1-(g_2^1-\bar{g}_2^1+c_2^1-s\,\omega_A^{1,2})$ is cohomologous to zero. Thus there is a component t_2^1 of T such that

$$\nabla(t_2^1)=\bar{g}\phi_2^1-(g_2^1-g_2^1+c_2^1+s\,\omega_A^{1,2}).$$

$\square$

We shall refer to the algorithm in the proof of the Transfer Theorem as the *Transfer Algorithm*. Since $\tilde{g}$ is a homology isomorphism whenever A is free (cf. Proposition 1) we have:

Corollary 1. *Let (A,d_A) be a free DGM, let (B,d_B,ω_B) be an A_∞-bialgebra, and let $g:A\to B$ be a chain map that is also a homology isomorphism. Then*

- *(i) (Existence) g induces an A_∞-bialgebra structure ω_A on A and extends to a map $G:A\Rightarrow B$ of A_∞-bialgebras.*
- *(ii) (Uniqueness) (ω_A,G) is unique up to isomorphism.*

Given a chain complex B of (not necessarily free) R-modules, there is always a chain complex of free R-modules (A, d_A) and a homology isomorphism $g : A \to B$. To see this, let $\left(RH : \cdots \to R_1H \to R_0H \xrightarrow{\rho} H, d\right)$ be a free R-module resolution of $H = H_*(B)$. Since R_0H is projective, there is a cycle-selecting homomorphism $g_0' : R_0H \to Z(B)$, which lifts the map ρ through the projection $Z(B) \to H$ and extends to a chain map $g_0 : (RH, 0) \to (B, d_B)$. If $RH : 0 \to R_1H \to R_0H \to H$ is a short R-module resolution of H, then g_0 extends to a homology isomorphism $g : (RH, d + h) \to (B, d_B)$ with $(A, d_A) = (RH, d)$. Otherwise, there is a perturbation h of d such that $g : (RH, d + h) \to (B, d_B)$ is a homology isomorphism with $(A, d_A) = (RH, d + h)$ (see [3], [17]). Thus an A_∞-structure on B always transfers to an A_∞-structure on $(RH, d + h)$ via Corollary 1, and we obtain our main result concerning the transfer of A_∞-structure to homology:

Theorem 2. *Let B be an A_∞-bialgebra with homology $H = H_*(B)$, let (RH, d) be a free R-module resolution of H, and let h be a perturbation of d such that $g : (RH, d + h) \to (B, d_B)$ is a homology isomorphism. Then*

- *(i) (Existence) g induces an A_∞-bialgebra structure ω_{RH} on RH and extends to a map $G : RH \Rightarrow B$ of A_∞-bialgebras.*
- *(ii) (Uniqueness) (ω_{RH}, G) is unique up to isomorphism.*

Remark 1. *Note that A_∞-bialgebra structures induced by the Transfer Algorithm are isomorphic for all choices of the map $g : (RH, d + h) \to (B, d_B)$, and we obtain an isomorphism class of A_∞-bialgebra structures on RH.*

Remark 2. *When $H = H_*(B)$ is a free module, we can set $RH = H$ and recover the classical results of Kadeishvili* [11]*, Markl* [15]*, and others, which transfer a DG (co)algebra structure to an A_∞-(co)algebra structure on homology. For an example of a DGA structure that cannot be transferred using classical techniques, but to which our method applies see* [26]*. Furthermore, any pair of A_∞-(co)algebra structures $\left\{\omega_H^{n,1}\right\}_{n\geq 1}$ and $\left\{\omega_H^{1,m}\right\}_{m\geq 1}$ on H induced by the same cycle-selecting map $g : H \to B$ extend to an A_∞-bialgebra structure $(H, \omega_H^{n,m})$, by the proof of Theorem 2.*

7. APPLICATIONS AND EXAMPLES

The applications and examples in this section apply the Transfer Algorithm given by the proof of Theorem 1. Three kinds of specialized A_∞-bialgebras $(A, \{\omega_m^n\})$ are relevant here:

1. $\omega_m^1 = 0$ for $m \geq 3$ (the A_∞-algebra substructure is trivial).
2. $\omega_m^n = 0$ for $m, n \geq 2$ (all higher order structure is concentrated in the A_∞-algebra and A_∞-coalgebra substructures).
3. Conditions (1) and (2) hold simultaneously.

Of these, A_∞-bialgebras of the first and third kind appear in the applications.

Structure relations defining A_∞-bialgebras of the second and third kind are expressed in terms of the S-U diagonal on associahedra Δ_K [20] and have especially nice form. Structure relations of the second kind were derived in [25]. Structure relations in an A_∞-bialgebra (A, ω) of the third kind with $\omega_1^1 = 0$, $\mu = \omega_2^1$ and $\psi^n = \omega_1^n$ are a special case of those derived in [25] and are given by the formula

$$\left\{\psi^n \mu = \mu^{\otimes n} \Psi^n\right\}_{n\geq 2}, \tag{7.1}$$

where the n-ary A_∞-coalgebra operation

$$\Psi^n = (\sigma_{n,2})_* \, \iota \, (\xi \otimes \xi) \, \Delta_K \left(e^{n-2} \right) : A \otimes A \to (A \otimes A)^{\otimes n}$$

is defined in terms of

- a map $\xi : C_*(K) \to Hom(A, TA)$ of operads sending the top dimensional cell $e^{n-2} \subseteq K_n$ to ψ^n,
- the canonical isomorphism
$$\iota : Hom \left(A, A^{\otimes n} \right)^{\otimes 2} \to Hom \left(A^{\otimes 2}, \left(A^{\otimes n} \right)^{\otimes 2} \right),$$
- and the induced isomorphism
$$(\sigma_{n,2})_* : Hom \left(A^{\otimes 2}, \left(A^{\otimes n} \right)^{\otimes 2} \right) \to Hom \left(A^{\otimes 2}, \left(A^{\otimes 2} \right)^{\otimes n} \right).$$

Structure relations defining a morphism $G = \{g^n\} : (A, \omega_A) \Rightarrow (B, \omega_B)$ between A_∞-bialgebras of the third kind are expressed in terms of the S-U diagonal on multiplihedra Δ_J [20] by the formula

$$\left\{ g^n \mu_A = \mu_B^{\otimes n} \mathbf{g}^n \right\}_{n \geq 1}, \tag{7.2}$$

where

$$\mathbf{g}^n = (\sigma_{n,2})_* \iota (\upsilon \otimes \upsilon) \Delta_J (e^{n-1}) : A \otimes A \to (B \otimes B)^{\otimes n},$$

and $\upsilon : C_*(J) \to Hom(A, TB)$ is a map of relative prematrads sending the top dimensional cell $e^{n-1} \subseteq J_n$ to g^n (the maps $\{\mathbf{g}^n\}$ define the tensor product morphism $G \otimes G : (A \otimes A, \Psi_{A \otimes A}) \Rightarrow (B \otimes B, \Psi_{B \otimes B})$).

Given a simply connected topological space X, consider the Moore loop space ΩX and the simplicial singular cochain complex $S^*(\Omega X; R)$. Under the hypotheses of the Transfer Theorem, the DG bialgebra structure of $S^*(\Omega X; R)$ transfers to an A_∞-bialgebra structure on $H^*(\Omega X; R)$. Our next two theorems apply this principle and identify some important A_∞-bialgebras of the third kind on loop space (co)homology.

Theorem 3. *If X is simply connected, $H^*(\Omega X; \mathbb{Q})$ admits an induced A_∞-bialgebra structure of the third kind.*

Proof. Let $\mathcal{A}_X$ be a free DG commutative algebra cochain model for X over $\mathbb{Q}$ (e.g., Sullivan's minimal or Halperin-Stasheff's filtered model); then $H^*(\mathcal{A}_X) \approx H^*(X; \mathbb{Q})$. The bar construction $(B = B\mathcal{A}_X, d_B, \Delta_B)$ with shuffle product is a cofree DG commutative Hopf algebra cochain model for ΩX, and $H = H^*(B, d_B)$ is a Hopf algebra with induced coproduct $\psi^2 = \omega_1^2$ and free graded commutative product $\mu = \omega_2^1$ (by a theorem of Hopf). Since H is a free commutative algebra, there is a multiplicative cocycle-selecting map $g_1^1 : H \to B$. Consequently, we may set $\omega_n^1 = 0$ for all $n \geq 3$ and $g_n^1 = 0$ for all $n \geq 2$ and obtain a trivial A_∞-algebra structure (H, μ) induced by g_1^1. There is an induced A_∞-coalgebra structure $(H, \psi^n)_{n \geq 2}$ and an A_∞-coalgebra map $G = \left\{ g^n \mid g^1 = g_1^1 \right\}_{n \geq 1} : H \Rightarrow B$ constructed as follows: For $n \geq 2$, assume ψ^n and g^{n-1} have been constructed, and apply the Transfer Algorithm to obtain candidates ω_1^{n+1} and g_1^n. Restrict ω_1^{n+1} to generators and let ψ^{n+1} be the free extension of ω_1^{n+1} to all of H using Formula 7.1. Similarly, restrict g_1^n to generators and let g^n be the free extension of g_1^n to all of H using Formula 7.2.

To complete the proof, we show that all other A_∞-bialgebra operations may be trivially chosen. Refer to the Transfer Algorithm and note that the Hopf relation

$\psi^2\mu = (\mu\otimes\mu)\,\sigma_{2,2}\left(\psi^2\otimes\psi^2\right)$ implies $\beta(\partial\mathfrak{f}_2^2)|_{C_1(JJ_{2,2}\setminus\operatorname{int}(KK_{2,2}\times 1))} = 0$. Thus we may choose $\omega_2^2 = g_2^2 = 0$ so that $\beta(\partial\mathfrak{f}_2^2) = \beta(\mathfrak{f}_2^2) = 0$. Inductively, assume that $\omega_2^{n-1} = g_2^{n-1} = 0$ for $n \geq 3$. Then $\beta(\partial\mathfrak{f}_2^n)|_{C_{n-1}(JJ_{n,2}\setminus\operatorname{int}(KK_{n,2}\times 1))} = 0$ and we may choose $\omega_2^n = 0$ and $g_2^n = 0$ so that $\beta(\partial\mathfrak{f}_2^n) = \beta(\mathfrak{f}_2^n) = 0$. Finally, for $m \geq 3$ set $\omega_m^n = 0$ and $g_m^n = 0$. Then $(H,\mu,\psi^n)_{n\geq 2}$ as an A_∞-bialgebra of the third kind with structure relations given by Formula 7.1, and G is a map of A_∞-bialgebras satisfying Formula 7.2. □

Note that the components of the A_∞-bialgebra map G in the proof of Theorem 3 are exactly the components of a map of underlying A_∞-coalgebras given by the Transfer Algorithm.

Let R be a PID and let X be a connected space such that $H_*(X;R)$ is torsion free. Then the Bott-Samelson Theorem [4] asserts that $H_*\left(\Omega\Sigma X;R\right)$ is isomorphic as an algebra to the tensor algebra $T^a\tilde{H}_*(X;R)$ generated by the reduced homology of X, and the adjoint $i : X \to \Omega\Sigma X$ of the identity $\mathbf{1} : \Sigma X \to \Sigma X$ induces the canonical inclusion $i_* : \tilde{H}_*\left(X;R\right) \hookrightarrow T^a\tilde{H}_*(X;R) \approx H_*\left(\Omega\Sigma X;R\right)$. Thus if $\{\psi^n\}_{n\geq 2}$ is an A_∞-coalgebra structure on $H_*(X;R)$, the tensor algebra $T^a\tilde{H}_*(X;R)$ admits a canonical A_∞-bialgebra structure of the third kind with respect to the free extension of each ψ^n via Formula 7.1.

Furthermore, the canonical inclusion $t : X \hookrightarrow \Omega\Sigma X$ induces a DG coalgebra map of simplicial singular chains $t_\# : S_*(X;R) \to S_*(\Omega\Sigma X;R)$, which extends to a homology isomorphism $t_\# : T^a\tilde{S}_*(X;R) \approx S_*(\Omega\Sigma X;R)$ of DG Hopf algebras. Thus the induced *Bott-Samelson Isomorphism* $t_* : T^a\tilde{H}_*\left(X;R\right) \approx H_*(\Omega\Sigma X;R)$ is an isomorphism of Hopf algebras ([10], [12]), and $T^a\tilde{S}_*(X;R)$ is a free DG Hopf algebra chain model for $\Omega\Sigma X$.

Theorem 4. *Let R be a PID and let X be a connected space such that $H_*(X;R)$ is torsion free.*

- *(i) Then $T^a\tilde{H}_*(X;R)$ admits an A_∞ -bialgebra structure of the third kind, which is trivial if and only if the A_∞-coalgebra structure of $H_*(X;R)$ is trivial.*
- *(ii) The Bott-Samelson Isomorphism $t_* : T^a\tilde{H}_*(X;R) \approx H_*(\Omega\Sigma X;R)$ extends to an isomorphism of A_∞-bialgebras.*

Proof. Since $H_*(X;R)$ is free as an R-module, we may choose a cycle-selecting map $\bar{g} = \bar{g}_1^1 : H_*(X,R) \to S_*(X;R)$ and apply the Transfer Algorithm to obtain an induced A_∞-coalgebra structure $\bar{\omega} = \{\bar{\omega}_1^n\}_{n\geq 2}$ on $H_*(X,R)$ and a corresponding map of A_∞-coalgebras $\bar{G} = \{\bar{g}_1^n\}_{n\geq 1} : H_*(X;R) \Rightarrow S_*(X;R)$. Let $H = T^a\tilde{H}_*(X;R)$, let $B = T^a\tilde{S}_*(X;R)$, and consider the free (multiplicative) extension $g = T\left(\bar{g}\right) : H \to B$. As in the proof of Theorem 3, use formulas 7.1 and 7.2 to freely extend $\bar{\omega}$ and $\bar{G}$ to families $\omega = \{\omega_1^n\}$ and $G = \{g_1^n \mid g_1^1 = g\}_{n\geq 1}$ defined on H, and choose all other A_∞-bialgebra operations to be zero. Then $\bar{\omega}$ lifts to an A_∞-bialgebra structure (H,ω,μ) of the third kind with free product μ, and $\bar{G}$ lifts to a map $G : H \Rightarrow B$ of A_∞-bialgebras. Furthermore, restricting ω to the multiplicative generators $H_*(X;R)$ recovers the A_∞-coalgebra operations on $H_*(X;R)$. Thus A_∞ -bialgebra structure of H is trivial if and only if the A_∞-coalgebra structure of $H_*(X;R)$ is trivial. Finally, since B is a free DG Hopf algebra chain model for $\Omega\Sigma X$, the Bott-Samelson Isomorphism t_* extends to an isomorphism of A_∞-bialgebras and identifies the A_∞-bialgebra structure of $H_*(\Omega\Sigma X;R)$ with (H,ω,μ). □

It is important to note that prior to this work, all known *rational* homology invariants of $\Omega\Sigma X$ are trivial for any space X. However, we now have the following:

Corollary 2. *A nontrivial A_∞-coalgebra structure on $H_*(X;\mathbb{Q})$ induces a nontrivial A_∞-bialgebra structure on $H_*(\Omega\Sigma X;\mathbb{Q})$. Thus the A_∞-bialgebra structure of $H_*(\Omega\Sigma X;\mathbb{Q})$ is a nontrivial rational homology invariant.*

Proof. First, $H = H_*(\Omega\Sigma X;\mathbb{Q})$ admits an induced A_∞-bialgebra structure of the third kind by Theorem 4, which is trivial if and only if the A_∞-coalgebra structure of $H_*(X;\mathbb{Q})$ is trivial. Second, the dual version of Theorem 3 imposes an induced A_∞-bialgebra structure on H whose A_∞-coalgebra substructure is trivial, and whose A_∞-algebra substructure is trivial if and only if the A_∞-coalgebra structure of $H_*(X;\mathbb{Q})$ is trivial. □

The two A_∞-bialgebras identified in the proof of Corollary 2 – one with trivial A_∞-coalgebra substructure and the other with trivial A_∞-algebra substructure – are in fact isomorphic, and represent the same isomorphism class of A_∞-bialgebra structures on $H_*(\Omega\Sigma X;\mathbb{Q})$ (cf. Remark 1). Indeed, choose a pair of isomorphisms for the two A_∞-(co)algebra substructures (their component in bidegree $(1,1)$ is $\mathbf{1} : H \to H$). Since $\omega_i^j = 0$ for $i, j \geq 2$, these isomorphisms clearly determine an isomorphism of A_∞-bialgebras.

Our next example exhibits an A_∞-bialgebra of the first but not the second kind. Given a 1-connected DGA (A, d_A), the bar construction of A, denoted by BA, is the cofree DGC $T^c\left(\downarrow \overline{A}\right)$ whose differential d and coproduct Δ are defined as follows: Let $\lfloor x_1|\cdots|x_n\rfloor$ denote the element $\downarrow x_1 \mid \cdots \mid\downarrow x_n \in BA$ and let e denote the unit $\lfloor\ \rfloor$. Then

$$d\lfloor x_1|\cdots|x_n\rfloor = \sum_{i=1}^{n} \pm \lfloor x_1|\cdots|d_A x_i|\cdots|x_n\rfloor + \sum_{i=1}^{n-1} \pm \lfloor x_1|\cdots|x_i x_{i+1}|\cdots|x_n\rfloor ;$$

$$\Delta\lfloor x_1|\cdots|x_n\rfloor = e \otimes \lfloor x_1|\cdots|x_n\rfloor + \lfloor x_1|\cdots|x_n\rfloor \otimes e + \sum_{i=1}^{n-1} \lfloor x_1|\cdots|x_i\rfloor \otimes \lfloor x_i|\cdots|x_n\rfloor .$$

Given an A_∞-coalgebra $(C, \Delta^n)_{n\geq 1}$, the tilde-cobar construction of C, denoted by $\tilde\Omega C$, is the free DGA $T^a\left(\uparrow \overline{C}\right)$ with differential $d_{\tilde\Omega A}$ given by extending $\sum_{i\geq 1}\Delta^n$ as a derivation. Let $\lceil x_1|\cdots|x_n\rceil$ denote $\uparrow x_1 \mid \cdots \mid\uparrow x_n \in \tilde\Omega H$.

Example 10. *Consider the DGA $A = \mathbb{Z}_2[a,b]/\left(a^4, ab\right)$ with $|a| = 3$, $|b| = 5$ and trivial differential. Define a homotopy Gerstenhaber algebra (HGA) structure $\{E_{p,q} : A^{\otimes p} \otimes A^{\otimes q} \to A\}_{p,q\geq 0;\ p+q>0}$ with $E_{p,q}$ acting trivially except $E_{1,0} = E_{0,1} = \mathbf{1}$ and $E_{1,1}(b;b) = a^3$ (cf.* [7], [12]*). Form the tensor coalgebra $BA \otimes BA$ with coproduct $\psi = \sigma_{2,2}(\Delta\otimes\Delta)$, and consider the induced map*

$$\phi = E'_{1,0} + E'_{0,1} + E'_{1,1} : BA \otimes BA \to A$$

of degree $+1$, which acts trivially except for $E'_{1,0}(\lfloor x\rfloor \otimes e) = E'_{0,1}(e \otimes \lfloor x\rfloor) = x$ for all $x \in A$, and $E'_{1,1}(\lfloor b\rfloor \otimes \lfloor b\rfloor) = a^3$. Since $E_{p,q}$ is an HGA structure, ϕ is a twisting cochain, which lifts to a chain map of DG coalgebras $\mu : BA \otimes BA \to BA$ defined by

$$\mu = \sum_{k=0}^{\infty} \downarrow^{\otimes k+1} \phi^{\otimes k+1} \bar\psi^{(k)},$$

where $\bar{\psi}^{(0)} = \mathbf{1}$, $\bar{\psi}^{(k)} = \left(\bar{\psi} \otimes \mathbf{1}^{\otimes k-1}\right) \cdots \left(\bar{\psi} \otimes \mathbf{1}\right) \bar{\psi}$ *for* $k > 0$, *and* $\bar{\psi}$ *is the reduced coproduct on* $BA \otimes BA$. *Then, for example,* $\mu\left(\lfloor b \rfloor \otimes \lfloor b \rfloor\right) = \left\lfloor a^3 \right\rfloor$ *and* $\mu\left(\lfloor b \rfloor \otimes \lfloor a|b \rfloor\right) = \left\lfloor a|a^3 \right\rfloor + \lfloor b|a|b \rfloor$. *It follows that* (BA, d, Δ, μ) *is a DG Hopf algebra. Let* μ_H *and* Δ_H *be the product and coproduct on* $H = H^*(BA)$ *induced by* μ *and* Δ; *then* (H, Δ_H, μ_H) *is a graded bialgebra. Let* $\alpha = \operatorname{cls} \lfloor a \rfloor$ *and* $z = \operatorname{cls} \left\lfloor a|a^3 \right\rfloor$ *in* H, *and note that* $\left\lfloor a^3 \right\rfloor = d \left\lfloor a|a^2 \right\rfloor$. *Let* $g : H \to BA$ *be a cycle-selecting map such that* $g\left(\operatorname{cls} \lfloor x_1 | \cdots | x_n \rfloor\right) = \lfloor x_1 | \cdots | x_n \rfloor$. *Then*

$$\bar{\Delta}_H(z) = \operatorname{cls} \bar{\Delta} \left\lfloor a|a^3 \right\rfloor = \operatorname{cls} \left\{ \lfloor a \rfloor \otimes \left\lfloor a^3 \right\rfloor \right\} = 0$$

so that

$$\left\{ \Delta g + (g \otimes g) \Delta_H \right\}(z) = \lfloor a \rfloor \otimes \left\lfloor a^3 \right\rfloor.$$

By the Transfer Theorem, we may choose a map $g^2 : H \to BA \otimes BA$ *such that* $g^2(z) = \lfloor a \rfloor \otimes \left\lfloor a|a^2 \right\rfloor$; *then*

$$\nabla g^2(z) = \left\{ \Delta g + (g \otimes g) \Delta_H \right\}(z).$$

Furthermore, note that

$$\left\{ \left(g^2 \otimes g + g \otimes g^2\right) \Delta_H + (\Delta \otimes \mathbf{1} + \mathbf{1} \otimes \Delta) g^2 \right\}(z) = \lfloor a \rfloor \otimes \lfloor a \rfloor \otimes \left\lfloor a^2 \right\rfloor.$$

Since $\left\lfloor a^2 \right\rfloor = d \lfloor a|a \rfloor$, *there is an* A_∞*-coalgebra operation* $\Delta_H^3 : H \to H^{\otimes 3}$ *and a map* $g^3 : H \to (BA)^{\otimes 3}$ *satisfying the general relation on* J_3 *such that* $\Delta_H^3(z) = 0$ *and* $g^3(z) = \lfloor a \rfloor \otimes \lfloor a \rfloor \otimes \lfloor a|a \rfloor$. *In fact, we may choose* Δ_H^3 *to be identically zero on* H *so that*

$$\nabla g^3 = (\Delta \otimes \mathbf{1} + \mathbf{1} \otimes \Delta) g^2 + \left(g^2 \otimes g + g \otimes g^2\right) \Delta_H.$$

Now the potentially non-vanishing terms in the image of J_4 *are*

$$\left(g^3 \otimes g + g^2 \otimes g^2 + g \otimes g^3\right) \Delta_H + (\Delta \otimes \mathbf{1} \otimes \mathbf{1} + \mathbf{1} \otimes \Delta \otimes \mathbf{1} + \mathbf{1} \otimes \mathbf{1} \otimes \Delta) g^3,$$

and evaluating at z *gives* $\lfloor a \rfloor \otimes \lfloor a \rfloor \otimes \lfloor a \rfloor \otimes \lfloor a \rfloor$. *Thus there is an* A_∞*-coalgebra operation* Δ_H^4 *and a map* $g^4 : H \to (BA)^{\otimes 4}$ *satisfying the general relation on* J_4 *such that* $\Delta_H^4(z) = \alpha \otimes \alpha \otimes \alpha \otimes \alpha$ *and* $g^4(z) = 0$. *Now recall that the induced* A_∞*-coalgebra structure on* $H \otimes H$ *is given by*

$$\begin{aligned}
\Delta_{H \otimes H} &= \sigma_{2,2} \left(\Delta_H \otimes \Delta_H\right) \\
\Delta_{H \otimes H}^4 &= \sigma_{4,2}[\Delta_H^4 \otimes (\mathbf{1}^{\otimes 2} \otimes \Delta_H)(\mathbf{1} \otimes \Delta_H)\Delta_H + (\Delta_H \otimes \mathbf{1}^{\otimes 2})(\Delta_H \otimes \mathbf{1})\Delta_H \otimes \Delta_H^4] \\
&\ \ \vdots
\end{aligned}$$

Let $\beta = \operatorname{cls} \lfloor b \rfloor$, $u = \operatorname{cls} \lfloor a|b \rfloor$, $v = \operatorname{cls} \lfloor b|a \rfloor$, *and* $w = \operatorname{cls} \lfloor b|a|b \rfloor$ *in* H, *and consider the induced map of tilde cobar constructions*

$$\widetilde{\mu_H} = \sum_{n \geq 1} (\uparrow \mu_H \downarrow)^{\otimes n} : \tilde{\Omega}(H \otimes H) \to \tilde{\Omega} H.$$

Then

$$\widetilde{\mu_H} \lceil \beta \otimes u \rceil = \lceil \mu_H (\beta \otimes u) \rceil = \lceil \operatorname{cls} \mu\left(\lfloor b \rfloor \otimes \lfloor a|b \rfloor\right) \rceil = \lceil z + w \rceil$$

so that

$$d_{\tilde{\Omega} H} \widetilde{\mu_H} \lceil \beta \otimes u \rceil = d_{\tilde{\Omega} H} \lceil z + w \rceil = \lceil \alpha|\alpha|\alpha|\alpha \rceil + \lceil \beta|u \rceil + \lceil v|\beta \rceil.$$

But on the other hand,

$$\begin{aligned} d_{\tilde{\Omega}(H\otimes H)}\left\lceil \beta\otimes u\right\rceil &= \left\lceil \Delta_{H\otimes H}\left(\beta\otimes u\right)\right\rceil \\ &= \left\lceil e\otimes u \mid \beta\otimes e\right\rceil + \left\lceil \beta\otimes e \mid e\otimes u\right\rceil \\ &\qquad + \left\lceil \beta\otimes\alpha \mid e\otimes\beta\right\rceil + \left\lceil e\otimes\alpha \mid \beta\otimes\beta\right\rceil \end{aligned}$$

so that

$$\widetilde{\mu_H} d_{\tilde{\Omega}(H\otimes H)}\left\lceil \beta\otimes u\right\rceil = \left\lceil \beta|u\right\rceil + \left\lceil v|\beta\right\rceil .$$

Although $\widetilde{\mu_H}$ fails to be a chain map, the Transfer Theorem implies there is a chain map $\widetilde{\mu_H}^2 : \tilde{\Omega}(H\otimes H)\to\tilde{\Omega}H$ such that $\widetilde{\mu_H}^2\left\lceil e\otimes\alpha \mid \beta\otimes\beta\right\rceil = \left\lceil \alpha|\alpha|\alpha|\alpha\right\rceil$, which can be realized by defining

$$\widetilde{\mu_H}^2 = \sum_{n\geq 1}\left(\uparrow\mu_H\downarrow + \uparrow^{\otimes 3}\omega_2^3\downarrow\right)^{\otimes n},$$

where $\omega_2^3\left(\beta\otimes\beta\right) = \alpha\otimes\alpha\otimes\alpha$. Indeed, to see that the required equality holds, note that $\mu_H\left(\beta\otimes\beta\right) = 0$ since $\left\lfloor a^3\right\rfloor = d\left\lfloor a|a^2\right\rfloor$. Thus there is a map $g_2 : H\otimes H\to BA$ such that $g_2\left(\beta\otimes\beta\right) = \left\lfloor a|a^2\right\rfloor$ and $\nabla g_2 = g\mu_H + \mu\left(g\otimes g\right)$. Furthermore, there is the following general relation on $JJ_{2,2}$:

$$\begin{aligned} \nabla g_2^2 &= \omega_{BA}^{2,2}\left(g\otimes g\right) + \left(\mu\otimes\mu\right)\sigma_{2,2}\left(\Delta g\otimes g^2 + g^2\otimes\left(g\otimes g\right)\Delta_H\right) + g^2\mu_H \\ &\quad + \left(\mu\left(g\otimes g\right)\otimes g_2 + g_2\otimes g\mu_H\right)\sigma_{2,2}\left(\Delta_H\otimes\Delta_H\right) + \Delta g_2 + \left(g\otimes g\right)\omega_2^2. \end{aligned} \tag{7.3}$$

The first expression on the right-hand side vanishes since BA has trivial higher order structure and the next two expressions vanish since $\mu_H\left(\beta\otimes\beta\right) = 0$ and $g^2(\beta) = 0$ (β is primitive). However,

$$\begin{aligned} &\left\{\left(\mu\left(g\otimes g\right)\otimes g_2 + g_2\otimes g\mu_H\right)\sigma_{2,2}\left(\Delta_H\otimes\Delta_H\right) + \Delta g_2\right\}\left(\beta\otimes\beta\right) \\ &\qquad = \bar{\Delta}\, g_2(\beta\otimes\beta) = \left\lfloor a\right\rfloor\otimes\left\lfloor a^2\right\rfloor . \end{aligned}$$

Since $d\left\lfloor a|a\right\rfloor = \left\lfloor a^2\right\rfloor$, there an operation $\omega_2^2 : H^{\otimes 2}\to H^{\otimes 2}$ and a map $g_2^2 : H^{\otimes 2}\to (BA)^{\otimes 2}$ satisfying relation (7.3) such that $\omega_2^2\left(\beta\otimes\beta\right) = 0$ and $g_2^2\left(\beta\otimes\beta\right) = \left\lfloor a\right\rfloor\otimes\left\lfloor a|a\right\rfloor$. Similarly, there is an operation $\omega_2^3 : H^{\otimes 2}\to H^{\otimes 3}$ and a map $g_2^3 : H^{\otimes 2}\to (BA)^{\otimes 3}$ satisfying the general relation on $JJ_{3,2}$ such that $\omega_2^3\left(\beta\otimes\beta\right) = \alpha\otimes\alpha\otimes\alpha$ and $g_2^3\left(\beta\otimes\beta\right) = 0$. Thus $\left(H,\mu_H,\Delta_H,\ \Delta_H^4,\omega_2^3,...\right)$ is an A_∞-bialgebra of the first kind.

One can think of the algebra A in Example 10 as the singular $\mathbb{Z}_2$-cohomology algebra of a space X with the Steenrod algebra $\mathcal{A}_2$ acting nontrivially via $Sq_1 b = a^3$ (recall that $Sq_1 : H^n\left(X;\mathbb{Z}_2\right)\to H^{2n-1}\left(X;\mathbb{Z}_2\right)$ is a homomorphism defined by $Sq_1\left[x\right] = \left[x\smile_1 x\right]$). Recall that a space X is $\mathbb{Z}_2$-*formal* if there exists a DGA B and cohomology isomorphisms $C^*(X;\mathbb{Z}_2)\leftarrow B\to H^*\left(X;\mathbb{Z}_2\right)$. Thus when X is $\mathbb{Z}_2$-formal, $H^*\left(BA\right)\approx H^*(\Omega X;\mathbb{Z}_2)$ as graded coalgebras. Now consider a $\mathbb{Z}_2$-formal space X whose cohomology $H^*\left(X;\mathbb{Z}_2\right)$ is generated multiplicatively by $\left\{a_1,\ldots,a_{n+1},b\right\}$, $n\geq 2$. Then Example 10 suggests the following conditions on X, which if satisfied, give rise to a nontrivial operation ω_2^n with $n\geq 2$, on the loop cohomology $H^*(\Omega X;\mathbb{Z}_2)$:

(1) $a_1 b = 0$;
(2) $a_1\cdots a_{n+1} = 0$;
(3) $a_{i_1}\cdots a_{i_k}\neq 0$ whenever $k\leq n$ and $i_p\neq i_q$ for all $p\neq q$;
(4) $Sq_1(b) = a_2\cdots a_{n+1}$.

To see this, consider the non-zero classes $\alpha_i = \operatorname{cls}\lfloor a_i \rfloor$, $\beta = \operatorname{cls}\lfloor b \rfloor$, $u = \operatorname{cls}\lfloor a_1|b \rfloor$, $w = \operatorname{cls}\lfloor b|a_1|b \rfloor$, and $z = \operatorname{cls}\lfloor a_1|a_2\cdots a_{n+1} \rfloor$ in $H = H^*(BA)$. Conditions (2) and (3) give rise to an induced A_∞-coalgebra structure $\left\{\Delta_H^k : H \to H^{\otimes k}\right\}$ such that $\Delta_H^k(z) = 0$ for $3 \leq k \leq n$ and $\Delta_H^{n+1}(z) = \alpha_1 \otimes \cdots \otimes \alpha_{n+1}$ with $g^k(z) = \lfloor a_1 \rfloor \otimes \cdots \otimes \lfloor a_{k-1} \rfloor \otimes \lfloor a_k|a_{k+1}\cdots a_{n+1} \rfloor$ for $2 \leq k \leq n$ and $g^{n+1}(z) = 0$. Next, condition (4) implies $\beta \smile u = w + z$, and we can define $\omega_2^k(\beta \otimes \beta) = 0$ for $2 \leq k < n$ and $\omega_2^n(\beta \otimes \beta) = \alpha_2 \otimes \cdots \otimes \alpha_{n+1}$ with $g_2^1(\beta \otimes \beta) = \lfloor a_2|a_3\cdots a_{n+1} \rfloor$, $g_2^k(\beta \otimes \beta) = \lfloor a_2 \rfloor \otimes \cdots \otimes \lfloor a_k \rfloor \otimes \lfloor a_{k+1}\,|a_{k+2}\cdots a_{n+1} \rfloor$ for $2 \leq k \leq n-1$ and $g_2^n(\beta \otimes \beta) = 0$. Indeed, the Transfer Theorem implies the existence of an A_∞-bialgebra structure in which ω_2^n satisfies the required structure relation on $JJ_{n,2}$. Note that the $\mathbb{Z}_2$-formality assumption is in fact superfluous here, as it is sufficient for α_i, β, and u to be non-zero.

Spaces X with $\mathbb{Z}_2$-cohomology satisfying conditions (1)-(4) abound.

Example 11. *Given an integer* $n \geq 2$, *choose positive integers* $r_1, \ldots, r_{n+1}$ *and* $m \geq 2$ *such that* $r_2 + \cdots + r_{n+1} = 4m-3$. *Consider the "thick bouquet"* $S^{r_1} \underline{\vee} \cdots \underline{\vee} S^{r_{n+1}}$, *i.e.,* $S^{r_1} \times \cdots \times S^{r_{n+1}}$ *with top dimensional cell removed, and generators* $\bar{a}_i \in H^{r_i}(S^{r_i};\mathbb{Z}_2)$. *Also consider the suspension of complex projective space* $\Sigma\mathbb{C}P^{2m-2}$ *with generators* $\bar{b} \in H^{2m-1}(\Sigma\mathbb{C}P^{2m-2};\mathbb{Z}_2)$ *and* $Sq_1\bar{b} \in H^{4m-3}(\Sigma\mathbb{C}P^{2m-2};\mathbb{Z}_2)$. *Let* $Y_n = S^{r_1} \underline{\vee} \cdots \underline{\vee} S^{r_{n+1}} \vee \Sigma\mathbb{C}P^{2m-2}$, *and choose a map* $f : Y_n \to K(\mathbb{Z}_2, 4m-3)$ *such that* $f^*(\iota_{4m-3}) = \bar{a}_2 \cdots \bar{a}_{n+1} + Sq_1\bar{b}$. *Finally, consider the pullback* $p : X_n \to Y_n$ *of the following path fibration:*

$$\begin{tikzcd}
K(\mathbb{Z}_2, 4m-4) \arrow[r] & X_n \arrow[r] \arrow[d, "p"'] & \mathcal{L}K(\mathbb{Z}_2, 4m-3) \arrow[d] \\
& Y_n \arrow[r, "f"] & K(\mathbb{Z}_2, 4m-3) \\
& \bar{a}_2 \cdots \bar{a}_{n+1} + Sq_1\bar{b} & \iota_{4m-3} \arrow[l, "f^*"]
\end{tikzcd}$$

Let $a_i = p^*(\bar{a}_i)$ *and* $b = p^*\left(\bar{b}\right)$; *then* $a_1, \ldots, a_{n+1}, b$ *are multiplicative generators of* $H^*(X_n;\mathbb{Z}_2)$ *satisfying conditions (1) - (4) above. We remark that one can also obtain a space* X_2' *with a non-trivial* ω_2^2 *on its loop cohomology by setting* $Y_2' = \left(S^2 \times S^3\right) \vee \Sigma\mathbb{C}P^2$ *in the construction above (see* [26] *for details).*

Finally, we note that the cohomology of Eilenberg-MacLane spaces and Lie groups fail to satisfy all of (1)-(4), and it would not be surprising to find that the operations ω_2^n vanish in their loop cohomologies for all $n \geq 2$. In the case of Eilenberg-MacLane spaces, recent work of Berciano and the second author seems to support this conjecture. Indeed, each tensor factor $A = E(v, 2n+1) \otimes \Gamma(w, 2np+2) \subset H_*(K(\mathbb{Z}, n);\mathbb{Z}_p)$, $n \geq 3$, and p an odd prime, is an A_∞-bialgebra of the third kind of the form $\left(A, \Delta^2, \Delta^p, \mu\right)$ (see [2]).

HGAs with nontrivial actions of the Steenrod algebra $\mathcal{A}_2$ were first considered by the first author in [18] and [19]. In general, the Steenrod $\smile_1$-cochain operation (together with the other higher cochain operations) induces a nontrivial HGA structure on $S^*(X)$. However, the failure of the differential to be a $\smile_1$-derivation prevents an immediate lifting of the HGA structure to cohomology. Nevertheless, such liftings are possible in certain situations, as we have seen in Example 10. Here is another such example.

Example 12. *Let* $g : S^{2n-2} \to S^n$ *be a map of spheres and let* $Y_{m,n} = S^m \times \left(e^{2n-1} \cup_g S^n\right)$. *Let* $*$ *be the wedge point of* $S^m \vee S^n \subset Y_{m,n}$, *let* $f : S^{2m-1} \to$

$S^m \times *$, and let $X_{m,n} = e^{2m} \cup_f Y_{m,n}$. Then $X_{m,n}$ is $\mathbb{Z}_2$-formal for each m and n (by a dimensional argument), and we may consider $A = H^*(X_{m,n};\mathbb{Z}_2)$ and $H = H^*(BA) \approx H^*(\Omega X_{m,n};\mathbb{Z}_2)$. Below we prove that:

- *(i) The A_∞-coalgebra structure of H is nontrivial if and only if the Hopf invariant $h(f) = 1$, in which case $m = 2, 4, 8$.*
- *(ii) If $h(f) = 1$, the A_∞-coalgebra structure on H extends to a nontrivial A_∞-bialgebra structure on H. Furthermore, let $a \in A^n$ and $c \in A^{2n-1}$ be multiplicative generators; then there is a perturbed multiplication φ on H if and only if $Sq_1 a = c$, in which case $n = 3, 5, 9$; otherwise φ is induced by the shuffle product on BA.*

Proof. Suppose $h(f) = 1$. Then A is generated multiplicatively by $a \in A^n$, $b \in A^m$, and $c \in A^{2n-1}$ subject to the relations

$$a^2 = c^2 = ac = ab^2 = b^2 c = b^3 = 0.$$

Let $\alpha = \operatorname{cls}\lfloor a\rfloor$, $\beta = \operatorname{cls}\lfloor b\rfloor$, $\gamma = \operatorname{cls}\lfloor c\rfloor$, and $z = \operatorname{cls}\lfloor b^2\rfloor \in H = H^*(BA)$. Given $x_i = \operatorname{cls}\lfloor u_i\rfloor \in H$ with $u_i u_{i+1} = 0$, let $x_1|\cdots|x_n = \operatorname{cls}\lfloor u_1|\cdots|u_n\rfloor$. Note that $x = \alpha|z = z|\alpha$ and $y = \gamma|z = z|\gamma$. Let Δ_H denote the coproduct in H induced by the cofree coproduct Δ in BA. Then x and y are primitive, and $\Delta_H(\alpha|z|\alpha) = e \otimes \alpha|z|\alpha + x \otimes \alpha + \alpha \otimes x + \alpha|z|\alpha \otimes e$. Define $g(x) = \lfloor a|b^2\rfloor$ and $g^2(x) = \lfloor a\rfloor \otimes \lfloor b|b\rfloor$; define $g(\alpha|z|\alpha) = \lfloor a|b^2|a\rfloor$ and $g^2(\alpha|z|\alpha) = \lfloor a\rfloor \otimes (\lfloor a|b|b\rfloor + \lfloor b|a|b\rfloor + \lfloor b|b|a\rfloor)$. There is an induced A_∞-coalgebra operation $\Delta_H^3 : H \to H^{\otimes 3}$, which vanishes except on elements of the form $\cdots|z|\cdots$, and may be defined on the elements x, y, and $\alpha|z|\alpha$ by

$$\begin{aligned}
&\Delta_H^3(x) = \alpha \otimes \beta \otimes \beta,\\
&\Delta_H^3(y) = \gamma \otimes \beta \otimes \beta, \text{ and}\\
&\Delta_H^3(\alpha|z|\alpha) = \alpha \otimes (\alpha|\beta + \beta|\alpha) \otimes \beta + \alpha \otimes \beta \otimes (\alpha|\beta + \beta|\alpha).
\end{aligned}$$

Then $\{\Delta_H, \Delta_H^3\}$ defines an A_∞-coalgebra structure on H. Furthermore, if $Sq_1 a = c$, which can only occur when $n = 3, 5, 9$, the induced HGA structure on A is determined by Sq_1 and induces a perturbation of the shuffle product $\mu : BA \otimes BA \to BA$ with $\mu(\lfloor a\rfloor \otimes \lfloor a\rfloor) = \lfloor c\rfloor$. The product μ lifts to a perturbed product φ on H such that

$$\varphi(\alpha \otimes \alpha|z) = \alpha|z|\alpha + \gamma|z,$$

and the A_∞-coalgebra structure $(H, \Delta_H, \Delta_H^3)$ extends to an A_∞-bialgebra structure $(H, \Delta_H, \Delta_H^3, \varphi)$ as in Example 10. On the other hand, if $Sq_1 a = 0$, then is induced by the shuffle product on BA and $\varphi(\alpha \otimes \alpha|z) = \alpha|z|\alpha$. Conversely, if $h(f) = 0$, then $b^2 = 0$ so that $\Delta_H^k = 0$, for all $k \geq 3$. □

We conclude with an investigation of the A_∞-bialgebra structure on the double cobar construction. To this end, we first prove a more general fact, which follows our next definition:

Definition 20. *Let (A, d, ψ, φ) be a free DG bialgebra, i.e., free as a DGA. An* ***acyclic cover of*** *A is a collection of acyclic DG submodules*

$$\mathcal{C}(A) = \{C^a \subseteq A \mid a \text{ is a monomial of } A\}$$

such that $\psi(C^a) \subseteq C^a \otimes C^a$ and $\varphi(C^a \otimes C^b) \subseteq C^{ab}$.

Proposition 2. *Let (A, d, ψ, φ) be a free DG bialgebra with acyclic cover $\mathcal{C}(A)$.*

- *(i) Then φ and ψ extend to an A_∞-bialgebra structure of the third kind.*

(ii) Let $(A', d', \psi', \varphi')$ be a free DG bialgebra with acyclic cover $\mathcal{C}(A')$, and let $f : A \to A'$ be a DGA map such that $f(C^a) \subseteq C^{f(a)}$ for all $C^a \in \mathcal{C}(A)$. Then f extends to a morphism of A_∞-bialgebras.

Proof. Define an A_∞-coalgebra structure as follows: Let $\psi^2 = \psi$; arbitrarily define ψ^3 on multiplicative generators and extend ψ^3 to decomposables via

$$\psi^3 \mu_A = \mu_A^{\otimes 3} \sigma_{3,2} \left(\psi^3 \otimes \psi^3 \right).$$

Inductively, if $\left\{ \psi^i \right\}_{i<n}$ have been constructed, arbitrarily define ψ^n on multiplicative generators and use (7.1) to extend ψ^n to decomposables. Since each ψ^n preserves $\mathcal{C}(A)$ by hypothesis, $\left\{ \varphi, \psi^2, \psi^3, \ldots \right\}$ is an A_∞-bialgebra structure of the third kind, as desired. The proof of part (ii) is similar. □

Given a space X, choose a base point $y \in X$. Let $\operatorname{Sing}^2 X$ denote the Eilenberg 2-subcomplex of $\operatorname{Sing} X$ and let $C_*(X) = C_*(\operatorname{Sing}^2 X)/C_{>0}(\operatorname{Sing} y)$. In [20] we constructed an explicit (non-coassociative) coproduct on the double-cobar construction $\Omega^2 C_*(X)$, which turns it into a DG bialgebra. Let $\boldsymbol{\Omega}^2$ denote the functor from the category of (2-reduced) simplicial sets to the category of permutahedral sets ([20], [13]) such that $\Omega^2 C_*(X) = C_*^{\diamondsuit}(\boldsymbol{\Omega}^2 \operatorname{Sing}^2 X)$, where $C_*^{\diamondsuit}(Y) = C_*\big(\operatorname{Sing}^M Y\big)/\langle \text{degeneracies} \rangle$, where $\operatorname{Sing}^M Y$ is the multipermutahedral singular complex of Y (see Definition 15 in [20]; cf. [1], [5]). Now consider the monoidal permutahedral set $\boldsymbol{\Omega}^2 \operatorname{Sing}^2 X$, and let V_* be its monoidal (non-degenerate) generators. For each $a \in V_n$, let

$$C^a = R \left\langle r\text{-faces of } a \mid 0 \leq r \leq n \right\rangle.$$

Then $\{C^a\}$ is an acyclic cover, and by Proposition 2 we immediately have:

Theorem 5. *The DG bialgebra structure on double cobar-construction $\Omega^2 C_*(X)$ extends to an A_∞-bialgebra of the third kind.*

Conjecture 1. *Given a 2-connected space X, the chain complex $C_*^{\diamondsuit}(\Omega^2 X)$ admits an A_∞-bialgebra structure extending the DG bialgebra structure constructed in [20]. Moreover, there exists a morphism of A_∞-bialgebras*

$$G = \{g_m^n\} : \Omega^2 C_*(X) \Rightarrow C_*^{\diamondsuit}(\Omega^2 X)$$

such that g_1^1 is a homology isomorphism.

References

[1] H. J. Baues, The Cobar Construction as a Hopf Algebra and the Lie Differential, Invent. Math., **132** (1998), 467–489.

[2] A. Berciano and R. Umble, Some Naturally Occurring Examples of A_∞-bialgebras, J. Pure and Appl. Algebra, **215** (2011), 1284–1291.

[3] N. Berikashvili, On the Differentials of Spectral Sequences (Russian), Proc. Tbilisi Mat. Inst., **51** (1976), 1–105.

[4] R. Bott and H. Samelson, On the Pontrjagin Product in Spaces of Paths, Comm. Math. Helv., **27** (1953), 320–337.

[5] G. Carlsson and R. J. Milgram, Stable Homotopy and Iterated Loop Spaces, Handbook of Algebraic Topology (Edited by I. M. James), North-Holland (1995), 505–583.

[6] M. Gerstenhaber, The Cohomology Structure of an Associative Ring. Ann. of Math., **78** (2) (1963), 267–288.

[7] M. Gerstenhaber and A. A. Voronov, Higher Operations on the Hochschild Complex, Functional Analysis and its Applications, **29** (1995), 1–5.

[8] V.K.A.M. Gugenheim, On a Perturbation Theory for the Homology of the Loop Space, J. Pure and Appl. Algebra, **25** (1982), 197–205.

[9] J. Huebschmann and T. Kadeishvili, Small Models for Chain Algebras, Math. Z., **207** (1991), 245–280.

[10] D. Husemoller, Fibre Bundles, third ed., Graduate Texts in Mathematics, no. 20, Springer-Verlag, New-York, 1994.

[11] T. Kadeishvili, On the Homology Theory of Fibre Spaces, Russian Math. Survey, **35** (1980), 131–138.

[12] T. Kadeishvili and S. Saneblidze, A Cubical Model of a Fibration, J. Pure and Appl. Algebra, **196** (2005), 203–228.

[13] ——————, The Twisted Cartesian Model for the Double Path Space Fibration, AT/0210224, preprint.

[14] J.-L. Loday and M. Ronco, Hopf Algebra of the Planar Binary Trees, *Adv. in Math.* **139,** No. 2 (1998), 293-309.

[15] M. Markl, Transfering A_∞ (Strongly Homotopy Associative) Structures, Rendiconti del Circolo Matematico di Palermo, Serie II, **79** (2006) 139–151.

[16] M. Markl, S. Shnider and J. Stasheff, "Operads in Algebra, Topology and Physics," Mathematical surveys and Monographs, **96** (2002).

[17] S. Saneblidze, On Derived Categories and Derived Functors, Extracta Mathematicae, **22** (2007), 315–324.

[18] ——————, The Hochschild Chain Complex of a Space is the Complex of the Hochschild Set, unpublished manuscript.

[19] ——————, Filtered Hirsch algebras, preprint (2011), arXiv:0707.2165.

[20] S. Saneblidze and R. Umble, Diagonals on the Permutahedra, Multiplihedra and Associahedra, *J. Homology, Homotopy and Appl.*, **6** (1) (2004), 363–411.

[21] ——————, The Biderivative and A_∞-bialgebras, *J. Homology, Homotopy and Appl.*, **7** (2) (2005), 161–177.

[22] ——————, Matrads, Biassociahedra, and A_∞-bialgebras, *J. Homology, Homotopy and Appl.*, **13** (1) (2011), 1–57.

[23] J. D. Stasheff, Homotopy Associativity of H-spaces I, II, Trans. A. M. S. **108** (1963), 275–312.

[24] ——————, "H-spaces from a Homotopy Point of View," SLNM 161, Springer, Berlin (1970).

[25] R. Umble, Structure Relations in Special A_∞-bialgebras, J. Math. Sci. **152** (3) (2008), 443–450.

[26] ——————, Non-operadic Operations on Loop Cohomology, J. Math. Sci., to appear; preprint (2012) arXiv:1108.5693.

A. Razmadze Mathematical Institute, Department of Geometry and Topology, M. Aleksidze St. 1, 0193 Tbilisi, Georgia

E-mail address: `sane@rmi.ge`

Department of Mathematics, Millersville University of Pennsylvania, Millersville, PA. 17551

E-mail address: `ron.umble@millersville.edu`